

his tool contains sample information, education and communications (IEC) materials on select components of the Minimum Initial Service Package (MISP) for reproductive health and family planning. The MISP is a coordinated set of priority interventions that should be implemented at the onset of every new emergency to prevent and manage the consequences of sexual violence, reduce HIV transmission, and prevent excess maternal and newborn morbidity and mortality. The MISP has been an international standard since 1997, and a Sphere standard since 2004.

The basic presentation of the IEC materials considers the varying demographic characteristics, literacy levels, and cultural acceptance of reproductive health among community members in humanitarian settings. In order to convey the key messages most effectively, pictorial images and simple language are used. Images are in black and white to keep them simple, easy to print, and to avoid the potentially offensive use of color.


The IEC literature generally supports culturally tailored images, including hairstyles, clothing and facial features. The "universal templates" have been developed for you to draw features that contextualize the images to your setting (such as hair and clothing). The "variation templates" are based on the "universal templates," and have been adapted to represent broad, but piloted ethnic features. All of the templates have a space for you to insert your agency logo, contact information and a small map. These templates have been developed so that you can have IEC materials readily available to share with the community once services have been established.

The interactive versions are available from http://www.iawg.net/iec/.


Template credits include: American College of Nurse-Midwives, Hesperian Foundation, World Health Organization, Meghan Fitzgerald and Stacey Patino.

Template A

Insert agency logo and map here.



For family planning information, counseling and contraceptives, talk to a health worker.




Template B

Insert agency logo and map here.




Spacing children well is good for proper care and well-being.




FAMILY PLANNING METHOD	PROTECTION FROM PREGNANCY	PROTECTION FROM SEXUALLY TRANSMITTED INFECTIONS
Barrier Methods Male Condom Female Condom	Good	Good
Hormonal Methods Birth control pills, injections Implants	Very Good *** ** ** ** ** ** ** ** **	None
LAM (Lactational Amenorrhea Method)	Good ★★	None
Withdrawal	Some	Some
Calendar Method	Good	None
IUD (Intra-Uterine Device)	Best ★★★★	None
Vasectomy/Tubal Ligation	Best ★★★★	None








FAMILY PLANNING METHOD	PROTECTION FROM PREGNANCY	PROTECTION FROM SEXUALLY TRANSMITTED INFECTIONS
Barrier Methods Male Condom	Good	Good
Female Condom		
Hormonal Methods Birth control pills, injections	Very Good	None
Implants	Best	
LAM (Lactational Amenorrhea Method)	Good	None
Withdrawal	Some	Some
Calendar Method	Good	None
IUD (Intra-Uterine Device)	Best	None
Vasectomy/Tubal Ligation	Best	None







How to use a male condom (Circumcised)



Check expiration date



Use a new condom every time you have sex



Place condom on tip of erect penis with rolled side out




Unroll condom all the way to base of penis



When finished, hold rim and tip of condom, and withdraw penis while it is still hard





Throw away used condom






How to use a male condom (Uncircumcised)







Template E

Insert agency logo and map here.

LAM


(Lactational Amenorrhea Method: **Exclusive Breastfeeding)**



The mother must not have menstruated since giving birth.



Breast milk must be the infant's only source of food.





What should you do if you think you may become pregnant after unprotected sex?

Insert agency logo and map here.















Go to a health worker as soon as possible for emergency contraceptive pills, within 5 days after unprotected sex.


If there are two pills, take both pills at the same time.






What to do after forced sex

Insert agency logo and map here.








Going to a health facility as soon as possible can help you get health care, prevent pregnancy and serious illness, and receive counseling.

Services are private and safe.





Template H








At the health center




- ◆ To prevent pregnancy, seek emergency contraception (EC) at the clinic within 5 days.
- To prevent sexually transmitted infections, ask for antibiotics.
- ◆ To prevent HIV, ask for post-exposure prophylaxis (PEP) within 3 days.
- ◆ To prevent tetanus, ask for tetanus toxoid within 1 week.
- ◆ To prevent hepatitis, ask for a vaccination within 2 weeks.




Template I


Preparing for Childbirth





Talk with family to plan for an emergency














Interchangeable Images

In some settings, the mode of transport, or health center symbol may be different. Please replace the images from templates F, G, H, I, J, K as appropriate.

