



Basic Emergency Obstetric and Newborn Care

in Humanitarian Settings: Select Signal Functions

Facilitator's Guide

Refresher Training Module for Health Care
Providers Implementing the MISp
Inter-agency Working Group on Reproductive
Health in Crises Training Partnership

Acknowledgements

This facilitator's guide has been made possible thanks to generous funding provided by USAID's Office of Foreign Disaster Assistance (OFDA). The project would not have been possible without ongoing collaboration among the membership of the Inter-agency Working Group on Reproductive Health in Crises (IAWG) through the efforts of the Training Partnership. The United Nations Population Fund (UNFPA), the Helping Mothers Survive Secretariat based at Jhpiego, and the Women's Refugee Commission have been of invaluable support to the production of this module.

Authors

This facilitator's guide was developed based on input from the IAWG membership. The materials were created based on World Health Organization (WHO) guidelines. The authors are grateful to Tomo Watanabe, who contributed substantially to drafting the training module content. Kristen Harker, Dr. Wilma Doedens, and Dr. Nguyen-Toan Tran also contributed technical input.

© 2017 Women's Refugee Commission

Women's Refugee Commission 15 West 37th Street, New York, NY 10018 212.551.3115

www.womensrefugeecommission.org

For more information on the Inter-agency Working Group on Reproductive Health in Crises and the IAWG-Training Partnership: www.iawg.net

Table of Contents:

Facilitator's Guide Overview	1
Sample Training Agenda	4
Preparation for a Training	8
Unit 1: Course Overview	12
Unit 2: Respectful Maternity Care in Emergencies	14
Unit 3: Oxytocic Drugs for Postpartum Hemorrhage	18
Unit 4: Manual Removal of the Placenta	25
Unit 5: Transport and Referral	30
Unit 6: Antibiotics for Postpartum Infection	34
Unit 7: Magnesium Sulfate for Severe Pre-eclampsia/Eclampsia	39
Unit 8: Rapid Assessment and Management	50
Unit 9: Newborn Resuscitation (Optional)	53
Unit 10: Partogram (Optional)	56
Unit 11: Closing	69
Sample Certificate of Attendance	70
References	71

Facilitator's Guide Overview

Introduction

The Minimum Initial Service Package (MISP) for Sexual and Reproductive Health (SRH) is a priority set of life-saving activities to be implemented at the onset of every humanitarian crisis. The MISP has five objectives¹:

1. Ensure the health sector/cluster identifies an organization and a SRH coordinator to lead and coordinate the implementation of the MISP.
2. Prevent and manage the consequences of sexual violence.
3. Reduce HIV transmission.
4. Prevent excess maternal and newborn morbidity and mortality.
5. Plan for comprehensive SRH services, integrated into primary health care as the situation permits.

Neglecting the MISP in humanitarian settings has serious consequences: preventable maternal and newborn deaths; sexual violence and subsequent trauma; sexually transmitted infections; unwanted pregnancies and unsafe abortions; and the possible spread of HIV.

Nurses, midwives, and doctors working in emergencies provide the clinical services needed to achieve objectives 2, 3, and 4 of the MISP. These services include clinical management of survivors of sexual violence and basic emergency obstetric care, while ensuring infection prevention practices at all times in all settings.

The IAWG designed a series of short clinical outreach refresher training modules to reinforce previously acquired knowledge and skills of health care staff that are tasked with providing these services. This module is one of four refresher training modules that together address all seven signal functions of basic emergency obstetric and newborn care (BEmONC). Information on the other refresher training modules – *Assisted Vaginal Delivery via Vacuum Extraction*, *Uterine Evacuation using Manual Vacuum Aspiration*, and the *Clinical Management of Sexual Violence Survivors* – is available through the IAWG Training Partnership.

Objective

The objective of the *Basic Emergency Obstetric and Newborn Care in Humanitarian Settings, Select Signal Functions: Facilitator's Guide* is to guide clinical trainers in conducting rapid face-to-face trainings on key components of BEmONC. The trainings are intended for crisis settings and to refresh health care providers' knowledge and skills.

Target Audience

This training module is designed for clinical service providers, including midwives, nurses or other mid-level providers, general practice physicians, and obstetricians/gynecologists who are currently attending or will attend deliveries in the acute phase of a humanitarian response.

¹ The MISP will be revised in 2017. Please be aware that this content is subject to modification.

Participant Prerequisites

Participants should already be able to:

- ◆ Demonstrate knowledge of the female reproductive system.
- ◆ Take a medical history and conduct a physical exam.
- ◆ Conduct spontaneous vaginal deliveries.

Participants should have received past training on:

- ◆ Performing spontaneous vaginal deliveries.
- ◆ Providing subsequent care for a mother and newborn.
- ◆ Recognizing and managing common complications of the peripartum period.
- ◆ Establishing intravenous access.

Assess Participants

It is important to assess the experience of participants prior to finalizing the participant roster. Only fully qualified clinical service providers who are currently attending normal and complicated deliveries are eligible for this short refresher course. To attend this course, participants must already be competent in performing normal vaginal deliveries.

If you find that there is a need for basic obstetric training for clinicians with no experience:

- ◆ Advise the participants' agency to plan for an extended BEmONC training.
- ◆ Inform the IAWG Training Partnership Coordinator (training@iawg.net).

Description of the Facilitator's Guide

This facilitator's guide includes presentation notes, case studies, and simulation activities. It provides information on the necessary skills for handling common obstetric emergencies at multiple levels of care, including postpartum infection, pre-eclampsia/eclampsia, and postpartum hemorrhage.

This training module is divided into ten units, which include the following elements:

1. **Time:** An estimate of how long it should take to complete the unit.
2. **Objectives:** Specific objectives to be met by the end of each unit.
3. **Materials:** Copies of all learner materials for distribution and answer keys for trainers.
4. **Instructions:** A list of the handouts, presentations, and other resources needed for each activity, followed by step-by-step guidance on how to facilitate interactive learning.

Trainers are highly encouraged to adapt the units to fit local training needs and objectives. This course is designed to be co-facilitated by a minimum of two trainers. The nature of the competency-based activities and concurrent simulation sessions requires more than one trainer to successfully implement the course. The recommended ratio is one facilitator per six learners.

In this guide, slide presentations are organized with a thumbnail image on the left and facilitation notes directly beside it on the right side of the page. The instructions walk the trainer through which points to emphasize and the interactive portions of the presentation.

Documentation and Certificates

Trainers should document participants' attendance and present certificates of attendance at the end of the course. Participants will also complete a multiple-choice pre-test and post-test on their knowledge in this area.

Competency Assessment

This module is a refresher training course for health service providers with existing skills. Trainers should assess both knowledge and skills to determine the competency of each participant. The pre-test and post-test serve as knowledge assessments. The checklists contained in this module can be used to assess skills during simulated practice. Trainers should identify and select clinical service providers deemed competent to perform the skills, and also identify learners who require further supportive supervision prior to performing the intervention.

Participant Evaluation

Trainers should conduct an informal process evaluation at the end of each workshop day to assess participants' satisfaction with the topics and activities. At the end of the course, participants should complete the final evaluation to provide feedback for further trainings.

Sample Agenda

This is an example of an agenda for this training module. Trainers may need to adjust the order of some content and the time allowed based on the setting and the experience level of the participants. There is an adaptable and printable version of the course agenda on the accompanying USB key and on the IAWG website.

UNIT	TIMING	CONTENT	OBJECTIVES <i>At the end of the unit, participants will be able to:</i>	METHODOLOGY
Day 1				
	8:00-8:30 (30 minutes)	Registration of participants <ul style="list-style-type: none"> Estimating blood loss stations Pre-test 		Learning activity
Introduction				
Unit 1	8:30-9:00 (30 minutes)	Welcome and introduction <ul style="list-style-type: none"> Icebreaker Expectations and ground rules 	<ul style="list-style-type: none"> reflect on their expectations of the training understand the objectives of the training agree on the ground rules of the training 	Presentation Discussion
Unit 2	9:00 – 9:20 (20 minutes)	Respectful maternity care in emergencies	<ul style="list-style-type: none"> discuss how basic emergency obstetric and newborn care (BEmONC) supports the implementation of the MISP in an emergency explain the concept of respectful maternity care 	Presentation Discussion
Oxytocic Drugs for Postpartum hemorrhage				
Unit 3	9:20-10:05 (45 min)	<ul style="list-style-type: none"> Preparing for delivery Identifying postpartum hemorrhage Management of postpartum hemorrhage 	<ul style="list-style-type: none"> identify the most common causes of postpartum hemorrhage following a normal vaginal delivery review management of bleeding after birth 	Role play Discussion Video
	10:05-10:35 (30 min)	<ul style="list-style-type: none"> Oxytocic drugs Appropriate use of oxytocic drugs 	<ul style="list-style-type: none"> accurately identify normal and abnormal postpartum blood loss verbalize supplies available in emergency kits to treat postpartum hemorrhage choose appropriate medications for managing postpartum hemorrhage 	Exercise Learning activity
	10:35-10:45 (10 min)	Referral to a higher level of care	identify appropriate treatment vs. referral for patients requiring emergency treatment	Documentation

UNIT	TIMING	CONTENT	OBJECTIVES <i>At the end of the unit, participants will be able to:</i>	METHODOLOGY
	10:45-11:30 (45 min)	<ul style="list-style-type: none"> Case study Group debriefing 	<ul style="list-style-type: none"> practice clinical decision-making for a patient with postpartum bleeding 	Small groups
	11:30-11:45 (45 min)	Break		
Manual Removal of the Placenta				
Unit 4	11:45-12:20 (35 min)	<ul style="list-style-type: none"> Review of third stage of labor Manual delivery of placenta 	<ul style="list-style-type: none"> recognize indications for manual removal of the placenta at multiple levels of care verbalize the procedure for manual removal of the placenta 	Presentation Video
	12:20-12:45 (25 min)	<ul style="list-style-type: none"> Active management of third stage of labor (AMTSL) 	<ul style="list-style-type: none"> demonstrate competence in AMTSL 	Discussion Learning activity
	12:45-13:45 (60 min)	Lunch		
	13:45-14:30 (45 min)	<ul style="list-style-type: none"> Group 1 – AMTSL Group 2 – Manual removal of placenta 	<ul style="list-style-type: none"> demonstrate competence in AMTSL demonstrate understanding of manual removal of placenta 	Skills practice Case study
	14:30-15:15 (45 min)	<ul style="list-style-type: none"> Group 2 – AMTSL Group 1 – Manual removal of placenta 	<ul style="list-style-type: none"> demonstrate competence in AMTSL demonstrate understanding of manual removal of placenta 	Skills practice Case study
	15:15-15:45 (30 min)	<ul style="list-style-type: none"> Group debriefing of skills practice 		Discussion
	15:45-16:00 (15 min)	Break		
	16:00-16:45 (45 min)	<ul style="list-style-type: none"> Transport and referral Intravenous (IV) insertion and fluid administration 	<ul style="list-style-type: none"> how to safely stabilize and prepare a patient for transport after postpartum hemorrhage review of IV insertion and fluid administration 	Presentation Skills practice
	16:45-18:00 (75 min)	<ul style="list-style-type: none"> Open skills practice 		

UNIT	TIMING	CONTENT	OBJECTIVES <i>At the end of the unit, participants will be able to:</i>	METHODOLOGY
Day 2				
Antibiotics for Postpartum Infection				
Unit 6	8:15-8:30 (15 min)	<ul style="list-style-type: none"> Review of previous day Introduction to the day Questions 		
	8:30-9:00 (30 min)	<ul style="list-style-type: none"> Identification and treatment of postpartum infections 	<ul style="list-style-type: none"> review assessment, diagnosis, treatment, and evaluation to care for a woman showing signs of infection in the postpartum period 	Presentation Discussion
	9:00-10:00 (60 min)	<ul style="list-style-type: none"> Case study Group debrief 	<ul style="list-style-type: none"> apply assessment, diagnosis, treatment, and evaluation for care for a patient showing signs of infection 	Learning activity Discussion
	10:00-10:15 (15 min)	Break		
Magnesium Sulfate for Severe Pre-eclampsia/Eclampsia				
Unit 7	10:15-11:00 (30 min)	<ul style="list-style-type: none"> Classifying hypertensive disorders of pregnancy Accurate blood pressure and urine assessment Assess danger signs 	<ul style="list-style-type: none"> understand the continuum of hypertensive disorders in pregnancy demonstrate an ability to accurately measure and record blood pressure demonstrate an ability to assess for severe pre-eclampsia and eclampsia in limited resource settings 	Role play Discussion Presentation
	11:00-12:00 (60 min)	<ul style="list-style-type: none"> Safe administration of magnesium sulfate Safe administration of antihypertensive medications 	<ul style="list-style-type: none"> demonstrate an ability to safely prepare magnesium sulfate for intramuscular (IM) and IV administration understand treatment protocols for antihypertensive medication administration 	Presentation Skills practice
	12:00-13:00 (60 min)	<ul style="list-style-type: none"> Clinical Simulation Debriefing 	<ul style="list-style-type: none"> diagnose and initiate treatment of a patient 	Simulation
	13:00-14:00 (60 min)	Lunch		

UNIT	TIMING	CONTENT	OBJECTIVES <i>At the end of the unit, participants will be able to:</i>	METHODOLOGY
Identification and Treatment of Shock				
Unit 8*	14:00-14:45 (45 min)	<ul style="list-style-type: none"> • Rapid assessment and management • Identification and treatment of shock 	<ul style="list-style-type: none"> • quickly identify and treat an emergency • treat the symptoms of shock 	
Summary and Closing				
Unit 11	14:45-15:00 (15 min)	<ul style="list-style-type: none"> • Review of available resources 	<ul style="list-style-type: none"> • understand further training and resources available for increasing skills 	
	15:00-15:30 (30 min)	<ul style="list-style-type: none"> • Ongoing education process and practice exercises • Clinical mentors 	<ul style="list-style-type: none"> • identify strong clinical leaders to champion ongoing practice in the health care facility • introduce the exercises and drill process for ongoing support and practice 	
	15:30-16:00 (30 min)	<ul style="list-style-type: none"> • Post-test • Course evaluation • Closing/Certificates 		

* **Unit 9: Newborn Resuscitation** – is available as an add-on to the schedule, as required by the context. This unit (45 minutes) can be added to the end of the second day, after the conclusion of Unit 8.

* **Unit 10: Partogram** – is available as an add-on to the schedule, as required by the context. This unit (45 minutes) can run concurrently with Unit 9, if using.

Preparation for the Training

Trainers should go through the tables below that outline the preparatory work that must be undertaken to successfully implement this course.

Course Materials List

This training module is designed to demonstrate and practice use of the supplies in the Inter-Agency Reproductive Health (RH) Kits. These supplies should be available in the setting where the participants are working at the time of the course.

The following is a complete list of all the medical equipment and course supplies that are needed for the successful implementation of the course. Verify that the medical equipment is already available at the training venue and make arrangements to bring any missing supplies with you.

Please note that any simulators provided by the IAWG Training Partnership are the responsibility of the facilitator and will need to be brought to and from the training location.

Course Materials Checklist

UNIT	MATERIALS	QTY	CHECK
All	A projector compatible to a computer with sound	1	
All	Flip charts with paper and markers	2	
All	Power supply	1	
All	Room for 15 participants with 2 tables, preferably at work site	1	
3,4,5,6	Handouts as indicated in the <i>Advanced Preparation Checklist</i>	1 per person	
All	Childbirth and neonatal simulators or local alternative	2	
3	Red dye or other appropriate simulated blood		
	• White towel with 600 cc simulated blood		
	• Gauze bandage with 100 cc simulated blood		
	• 300 cc simulated blood as liquid in a basin		
	• 500 cc red fruit jam		
	Simulated medications		
3	• Oxytocin, misoprostol, and ergometrine	2-3	
4	• Magnesium sulfate, calcium gluconate, 2% lignocaine	2-3	
5	• Clindamycin and gentamicin	2-3	
All	Normal delivery supplies		
	• Gloves – sterile and non-sterile	1-2 boxes	
	• Blood pressure cuff (sphygmomanometer)	2-3	
	• IV infusion start equipment	10	
	• Syringes and needles	10	

UNIT	MATERIALS	QTY	CHECK
	• Catheter for bladder	2	
	• Adult and newborn bag/mask	2	
	• Reflex hammer	2	
	• Bin for placenta	2	
	• Speculum	2	
	• Sponge forceps	2	
	• Needle holder	2	
	• Scissors	2	
	• Suture	2	
	• Melon or other local fruit	5-6	
	• Sharps container	2	

Note to Trainers: In crisis settings, refresher courses typically take place on-site. Be sure to bring all supplies to the training, including paper certificates, copies of handouts, and additional resources in paper format. Bring paper copies of slides and extra flip chart paper as back up.

Advance Preparation Checklist

Use the following checklist to ensure that all the necessary materials are prepared in the weeks before delivering the training:

ITEM TO PREPARE	UNIT	COMPLETE
Check projector for image and sound quality of videos and presentations	General	
Review all slide presentations and hide any optional slides if not applicable to the context	General	
Review the instructions and answer keys for all activities	General	
Check that all course materials are available and ready to use	General	
Ensure access to childbirth simulator and comfort with all functionalities	General	
Flipchart sheets on: Ground rules, parking lot, Icebreaker (if adapting)	General	
Gather all materials listed for the course	General	
Print one copy of the following hand-outs for each participant:		
Course agenda	1	
Pre-test	1	
Post-test	9	
Certificate of attendance	9	
Course evaluation	9	
Familiarize yourself with slides on the formulation of magnesium sulfate and how to explain dosage concentrations	7	
Set up blood estimation exercise materials	1 and 3	

Further Reading:

- ◆ Inter-agency Working Group on Reproductive Health in Crises, *Inter-agency Field Manual for Reproductive Health in Crises*, Ch. 6, Maternal and Newborn Health. Geneva. 2010.
- ◆ World Health Organization, *Integrated Management of Pregnancy and Childbirth*. Geneva. 2015.
- ◆ UNFPA, *Inter-agency Reproductive Health Kits for Crisis Situations*. 5th Ed. New York. 2011.

These and other publications can be downloaded at www.iawg.net or by contacting training@iawg.net.

Accompanying USB

The accompanying USB key includes several resources to be used in the workshop activities, including all handouts and slide presentations. The key also includes sample certificates of completion and competence, which trainers can personalize.

Materials are also available on the Training Partnership Initiative page of the IAWG website : iawg.net/tpi-home/. Please see the "Resources" tab for this and other trainings on the clinical components of the MISP.



Alert: Important recommendations for before and after the training

This module is designed to be a 2 or 3-day clinical refresher training for already trained health care service providers in humanitarian contexts. The IAWG Training Partnership Initiative conducted research about the barriers and facilitators to implementing such trainings in crises settings, and has the following recommendations **for trainers and program managers**:

- ◆ Before the training:
 - ❖ Limit the selection of participants to those who meet the specified pre-requisite qualifications in the facilitator's guide.
 - ❖ Assess qualified participants' learning needs to prepare for how best to address knowledge gaps. To do so, use the pre-test included in the *Select Signal Functions of Basic Emergency Obstetric and Newborn Care* module or interview participants.
 - ❖ Based on identified trainees' needs, provide additional training resources to expand the training as needed.
 - ❖ Ensure that participants become certified in the MISP Distance Learning Module (available at <http://iawg.net/minimum-initial-service-package/>) as a foundation prior to the training, if possible.
- ◆ After the training:
 - ❖ Discuss and possibly organize a log book and a calendar of opportunities for trainees to practice their skills at their institution.
 - ❖ Regularly schedule ongoing supportive supervision for the providers as soon as the security situation allows.

For more information, please see the following guidance note: *Sexual and Reproductive Health Clinical Outreach Refresher Trainings (S-CORTs): Operational considerations and programmatic guidance for SRH trainers, program managers, and coordinators*.

Characteristics of an Effective Training

The following recommendations are necessary to ensure the effective transfer of information during adult learning:

- ❖ Clearly communicate the purpose of the training to both trainers and learners;
- ❖ State exactly what learners are expected to do at the end of the course;
- ❖ Use training methods that build on participants' existing skills and experience, enabling them to meet the objectives;
- ❖ Present new knowledge and skills in a relevant context;
- ❖ Actively engage learners in the process;
- ❖ Use an effective mix of training methods to meet the needs of different learning styles;
- ❖ Offer learners the opportunity to practice applying new knowledge and skills;
- ❖ Provide learners with constructive feedback on their performance;
- ❖ Allow enough time for learners to meet the objectives of the training; and
- ❖ Offer trainers and learners the opportunity to evaluate the course, measuring the extent to which trainers and learners met the training objectives, and accept feedback from learners to make improvements to the course.

From Ipas, Women-Centered Postabortion Care: Reference Manual, Second edition. Chapel Hill. 2013.

Sensitivity and Flexibility in a Crisis Setting

Sensitivity and flexibility are crucial in a crisis setting. When planning this training, trainers should keep the following notes in mind:

- ◆ Minimize the time providers spend away from their duty stations.
- ◆ Be sensitive to the long hours and double shifts health care providers may be working.
- ◆ Remember that some participants may have long travel times.
- ◆ Be prepared for participants with a range of abilities and experiences – some participants may be very new to the setting.
- ◆ Be aware that participants will most likely include a mix of nurses, midwives, doctors, and clinical officers.
- ◆ Be sensitive to the emotional needs of health care providers/participants in a crisis setting.
- ◆ Remember that some experienced providers may be inexperienced in providing care during a crisis.

Unit 1: Course Overview

Time

30 minutes

Objectives

By the end of this unit, participants should be able to:

- ◆ Reflect on their expectations of the training.
- ◆ Understand the objectives of the training.
- ◆ Agree on the ground rules of the training.

Materials

Prepare the following in advance:

- ◆ Two flip charts: one titled "Ground Rules" and one titled "Parking Lot"
- ◆ *Course Agenda* and *Pre-test*, one each per participant
- ◆ Markers in various colors, three or four per table
- ◆ Post-it notes in various colors, three or four per participant
- ◆ Blood estimation materials
 - ❖ A dyed white towel or culturally appropriate cloth (600 cc simulated blood - red dye or an available alternative)
 - ❖ A dyed gauze bandage (100 cc simulated blood - red dye or an available alternative)
 - ❖ Red liquid in a basin (300 cc simulated blood - red dye or an available alternative)
 - ❖ Red fruit jam in a container to represent a blood clot (500 cc)

Instructions

Place the *Course Agenda*, markers, and Post-it notes on the tables for the participants prior to beginning the unit. There may be some formal opening requests from national dignitaries. Please try to accommodate such requests in the shortest time possible.

During this session, participants will be introduced to each other and the facilitators. Participants will conduct the blood estimations exercise below and complete and return the *Pre-test*.

Blood Estimation Exercise

Adapted with permission from Jhpiego. Helping Mothers Survive – Bleeding after birth. 2015.

Before the training begins, set up four stations showing blood loss in different amounts and ways with the materials outlined above, and label them A, B, C, and D. As the participants become settled, invite them to walk through the blood estimation exercise and record their estimates of how much blood they think was lost in each of the stations. The estimates will be discussed in Unit 3.

After all participants record their estimates, start the training and work through the rest of the material outlined in this unit.

Housekeeping

Tell participants where the restrooms are and encourage them to leave the training room quietly, if needed. Mention that there will be morning and afternoon breaks with lunch in between and where these will take place.

Ground Rules

Present a flipchart sheet titled "Ground Rules":

- ◆ Explain that ground rules are mutually agreed-upon guidelines to help the group work together, create a safe and respectful learning environment, and accomplish tasks efficiently.
- ◆ Ask participants to suggest ground rules. Write their suggestions on the flip chart.
- ◆ Possible ground rules may include participating, listening respectfully, speaking one at a time, turning off cell phones and pagers, and maintaining confidentiality.

Expectations and Agenda

Ask participants to write one to three expectations of what they hope to learn during the course on Post-it notes. Tell participants to use one Post-it note for each expectation. Have the participants put the notes on the wall or a flip chart at the front of the room. Review and group the expectations into categories. Briefly go through the agenda. Explain the objectives of the training and which expectations can and cannot be addressed.

Parking Lot

Present the flip chart titled "Parking Lot." Explain that questions that arise during the course that cannot be immediately addressed will be put in the "Parking Lot." Throughout the course, questions will be referred to and addressed when they are most relevant.

Knowledge Pre-test

Distribute the *Pre-test* (available on the USB key and IAWG website) and tell participants that they have about 15 minutes to complete the questions. Ask them to write their names at the top of the document, and collect all tests at the end of the allotted time.

Presentation



Unit 1: Course overview

Basic emergency obstetric and newborn care in humanitarian settings

IAWG INTER-AGENCY WORKING GROUP ON REPRODUCTIVE HEALTH IN CRISES

Unit 2: Respectful Maternity Care in Emergencies

Time

20 minutes

Objectives

By the end of this unit, participants will be able to:

- ◆ Discuss how basic emergency obstetric and newborn care supports the implementation of the MISP in an emergency.
- ◆ Explain the concept of respectful maternity care.

Materials

- ◆ Projector and computer with sound
- ◆ Slide presentation
- ◆ *Respectful Maternity Care Charter* handout, one per participant

Instructions

Follow the **slide presentation** and emphasize the notes listed below the slides. After reviewing the last slide about the right of childbearing women to respectful maternity care, **facilitate a short discussion** by asking the participants how they can demonstrate respect for the women they serve. Hand out the *Respectful Maternity Care Charter* to aid this discussion.

(The following responses are taken with permission from Jhpiego. Helping Mothers Survive – Bleeding after birth. 2015.)

Appropriate responses include:

- ◆ Introduce yourself by name and smile.
- ◆ Look at the woman when speaking to her.
- ◆ Use simple, clear language and explain everything you are going to do.
- ◆ Speak calmly.
- ◆ Pay attention when the woman speaks and listen to what she says.
- ◆ Include women and their families in discussions about care.
- ◆ Always explain procedures and get the woman's consent before you begin.

Next, ask the participants:

- ◆ Have you ever had a patient refuse care?
- ◆ Was this woman handled with respect?
- ◆ In the future, would you do anything differently in a similar situation?

End by emphasizing that all women deserve respect, including women affected by humanitarian emergencies.

Presentation



Unit 2: Respectful maternity care in emergencies

Basic emergency obstetric and newborn care in humanitarian settings



Background


- In a displaced population, approx. 4% of the total population will be pregnant at any given time
- Maternal deaths: Estimated 289,000 in 2013
- Lifetime risk of maternal death:
 - Sub-Saharan Africa: 1 in 38
 - Oceania: 1 in 140
 - Southern Asia: 1 in 200
- Causes
 - Severe bleeding
 - Infection
 - Eclampsia
 - Obstructed labor

61% of maternal deaths worldwide occur in fragile states, many of them affected by conflict and recurring natural disasters



The most common life-threatening obstetric complications are severe bleeding, infection, eclampsia, and obstructed labor. There is a separate clinical refresher training module devoted to obstructed labor. This module focuses on the skills related to managing severe bleeding, infection, and eclampsia.

Minimum Initial Service Package (MISP) for Reproductive Health



Review the MISP and explain that objective 4 includes the basic and comprehensive emergency obstetric care signal functions that are needed to prevent excess maternal and newborn morbidity and mortality.

Prevent excess maternal and newborn morbidity and mortality

- Emergency obstetric and newborn care services are available, accessible, and utilized
- 24/7 referral system established
- Clean delivery kits provided to birth attendants and visibly pregnant women



The detailed version of Objective 4 is further described below:

- ◆ Ensure availability, accessibility, and utilization of delivery, essential newborn care, and emergency obstetric and newborn care (EmONC) services:

At health facilities: This includes skilled birth attendants and supplies for normal births, essential newborn care, and management of basic obstetric and newborn complications (basic EmONC).

At referral hospitals: This includes all the health facility activities plus skilled medical staff and supplies for the management of comprehensive obstetric and newborn emergencies (comprehensive EmONC).

- ◆ Establish a referral system to facilitate transport and communication from the community to the health center, and between the health center and the hospital.
- ◆ Ensure availability of a package of supplies and commodities for clean delivery and newborn care, where access to a health facility is not possible.

Signal functions of basic emergency obstetric and newborn care

BASIC EmONC

1. Antibiotics IV/IM
2. Oxytocic drugs IV/IM
3. Anticonvulsants IV/IM
4. Manual removal of placenta
5. Manual vacuum aspiration of retained products of conception
6. Vacuum extraction
7. Newborn resuscitation

COMPREHENSIVE EmONC

7. Surgery, including cesarean section
8. Blood transfusion

This slide describes the signal functions of basic emergency obstetric care that should be readily available at community-level health facilities (1 facility per 30,000 people). The signal functions are the minimal competencies for health care providers when providing obstetric care.

Signal functions of basic emergency obstetric and newborn care

BASIC EmONC

1. Antibiotics IV/IM
2. Oxytocic drugs IV/IM
3. Anticonvulsants IV/IM
4. Manual removal of placenta
5. Manual vacuum aspiration of retained products of conception
6. Vacuum extraction
7. Newborn resuscitation

COMPREHENSIVE EmONC

7. Surgery, including cesarean section
8. Blood transfusion

Basic EmONC signal functions are a set of life saving services used in the treatment of acute obstetric complications. This outreach training focuses on signal functions one through four, to address the leading causes of maternal morbidity and mortality. Signal functions five, six, and seven are addressed in other refresher training modules.

Respectful maternity care

- Freedom from harm and ill treatment
- Right to informed consent and refusal
- Confidentiality and privacy
- Dignity and respect
- Equal care
- Timely health care
- Autonomy and freedom from coercion



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN EMERGENCIES

Every woman regardless of ethnic background, culture, social standing, religion, education level, and marital or economic status deserves respectful maternity care.

Ask the participants: How does respectful care save lives?

Answer: Women may not seek care if they expect they will not be treated well.

Unit 3: Oxytocic Drugs for Postpartum Hemorrhage

Time

90 minutes

Objectives

By the end of this unit, participants will be able to:

- ◆ Accurately identify normal and abnormal postpartum blood loss.
- ◆ Identify and manage the most common causes of postpartum hemorrhage following a normal vaginal delivery.
- ◆ Utilize supplies available in the Inter-Agency Reproductive Health Kits to treat postpartum hemorrhage.
- ◆ Identify and appropriately refer patients requiring a higher level of care.

Materials

- ◆ Video – postpartum hemorrhage (on the USB key and IAWG website)
- ◆ Normal delivery supplies (see the Course Materials Checklist)
- ◆ Childbirth simulator
- ◆ Simulated medications – oxytocin, misoprostol, ergometrine
- ◆ Intravenous (IV) fluids
- ◆ *Case Study* handout, one per participant

Instructions

This unit introduces a woman presenting to the health care facility with an imminent vaginal delivery. Explain that the woman is actively laboring and arrives close to delivery. **Role play the woman who is actively laboring** using a simulator or other available supplies. Have the second facilitator follow the steps below to model appropriate respectful maternity care and the steps of a normal vaginal delivery followed by a postpartum hemorrhage. If facilitating alone (it is highly recommended that two facilitators are present), then elicit help from the learners. This activity will give you an idea of how learners usually practice and what to emphasize during the training.

Role Play Checklist

Adapted with permission from Jhpiego. Helping Mothers Survive: Bleeding after Birth. 2015.

- ☐ Deliver baby onto mother's stomach.
- ☐ Dry baby thoroughly and assess for crying or breathing; cover with a dry cloth and place on mother's chest.
- ☐ Check for a second baby. If there is none, then continue with third stage care while continuing to observe the baby.
- ☐ Give a uterotonic (oxytocin or misoprostol) within one minute of delivery.
- ☐ While awaiting the placenta, remove the first pair of gloves – if double-gloved – or change gloves and clamp and cut the cord between 1-3 minutes after birth.
- ☐ Perform controlled cord traction during contractions.
- ☐ Feel the uterus once the placenta delivers and massage if soft.
- ☐ Stop the role play and address the learners indicating the mother has too much bleeding.

After the role play, guide a discussion using the following questions:

- ◆ Have you ever seen someone die from bleeding after birth?
- ◆ What happened?
- ◆ Was there anything else that could have been done if the mother had been somewhere else?

Note to Trainers: This refresher training assumes a baseline level of familiarity with management of postpartum hemorrhage and management of normal vaginal delivery. If the need for further training on normal childbirth is identified, then give appropriate feedback to the training coordinator.

Introduce the video (22 minutes) as a refresher to the steps of managing and treating postpartum hemorrhage. Show the video and place any questions after the video into the parking lot.

Review the estimates from the blood loss estimation exercise at the beginning of the day when you reach the slide about the amount of normal blood loss. Discuss participants' estimates and how easy it can be to underestimate blood loss.

Notes for the remainder of the slides are listed underneath each slide in the presentation.

At the end of the presentation, hand out the **case study** (as listed below) and instruct the participants to work in small groups to answer the questions. Come back together after 30 minutes. Review the answers to the case study and give feedback on general impressions of the skills checklist to the group. **Ask the participants** what they learned from the case study.

Case Study

Mrs. F is a 20-year-old para 1 who has been brought to the health center by the local traditional birth attendant (TBA) because she has been bleeding heavily since childbirth at home 2 hours ago. The TBA reports that the birth was a spontaneous vaginal delivery of a full-term newborn. Mrs. F and the TBA report that the duration of labor was 12 hours, the birth was normal, and the placenta was delivered 20 minutes after the birth of the newborn.

Assessment (history, physical examination, screening procedures/laboratory tests)

1. What will you include in your initial assessment of Mrs. F? Why?
 - ▼ Vital signs
 - ▼ Uterine tone
 - ▼ Amount of vaginal bleeding
 - ▼ Status of placenta
 - ▼ Medications or herbs given
2. What particular aspects of Mrs. F's physical examination will help you make a diagnosis or identify her problems/needs? Why?
 - ▼ Fundal tone
 - ▼ Amount vaginal bleeding
 - ▼ Presence of vaginal trauma
 - ▼ Presence of symptoms of shock

Diagnosis (identification of problems/needs)

You have completed your rapid assessment of Mrs. F, and your main findings include the following:

History

- ▼ The TBA says that she thinks the placenta and membranes were delivered without difficulty and were complete.

Physical examination

- ▼ Mrs. F's temperature is 36.8°C, her pulse rate is 108 per minute, her blood pressure is 80/60 and her respirations are 24 per minute. She is pale and sweating. Her uterus is soft and does not contract with fundal massage. She has heavy, bright red vaginal bleeding. On inspection, there is no evidence of perineal, vaginal, or cervical tears.

3. Based on these findings, what is Mrs. F's diagnosis (problem/need)? Why?

- ▼ Postpartum hemorrhage with uterine atony
- ▼ Shock

Care provision (planning and intervention)

4. Based on your diagnosis (problem/need identification), what is your plan of care for Mrs. F? Why?

- ▼ IV fluids
- ▼ Stabilize and refer
- ▼ Oxytocin and/or misoprostol
- ▼ Attempt manual removal of the placenta



Unit 3: Oxytocic drugs for postpartum hemorrhage (PPH)

Basic emergency obstetric and newborn care in humanitarian settings



Start this segment with the **role play** of a normal childbirth followed by a postpartum hemorrhage (PPH). Stop the role play when excessive bleeding with a soft uterus is identified.

Watch the PPH video (22 minutes)



Watch the video as a refresher to the steps of managing and treating postpartum hemorrhage.

If participants have questions, place them in the "parking lot" for now.

Be prepared at every birth


- Assume anemia and malnourishment
- Call for help quickly
- Know your supplies
 - How to start an IV and give fluids
 - Medication storage and administration




Women who have anemia or are malnourished have less reserves to cope with blood loss following childbirth. Be prepared to manage bleeding at every birth by knowing who to call for help and acting quickly, and having supplies available – IV, medication, and fluids.

Is bleeding normal?

350 mL



500 mL



IAWIG INTER-AGENCY WORKING GROUP ON REPRODUCTIVE HEALTH IN CRISES

All women are at risk for bleeding after birth. Bleeding can happen as a quick gush or be a slow, constant stream of blood. Both are dangerous. Losing more than 500 mL of blood is considered a hemorrhage. **Ask the participants to share their estimations for blood loss from the stations that were set up at the start of the training.** After hearing the responses, give the right answers. Explain that it is easy to underestimate blood loss and that it is important to act quickly to stop bleeding after birth.

Causes

Remember the four Ts:

- Tone
- Trauma
- Tissue
- Thrombin

Uterine atony is the most common cause of PPH

IAWIG INTER-AGENCY WORKING GROUP ON REPRODUCTIVE HEALTH IN CRISES

The priority in postpartum hemorrhage is to find the cause of the bleeding, stop the bleeding, and stabilize the patient. This refresher training emphasizes emergency management of uterine atony (tone) and retained placenta (tissue). Failure of the uterus to contract (atonic uterus) is the most common cause of bleeding after childbirth. After the placenta separates, the contractions of the uterus occlude the blood vessels supplying the placenta and prevent excessive blood loss.

If the uterus is hard and the mother is still bleeding, tears are likely even if they cannot be seen. Very rarely, clotting disorders also lead to excessive bleeding after childbirth.

Massage uterus

- Uterus bleeds excessively until it contracts
- Check uterine tone and massage if soft
- A contracted uterus feels hard and small like a grapefruit or coconut

IAWIG INTER-AGENCY WORKING GROUP ON REPRODUCTIVE HEALTH IN CRISES

Most bleeding after birth is caused by a uterus that will not get or stay hard (uterine atony) after the birth of the placenta. Blood vessels continue to pump blood into the uterus if the uterus is not hard. Contributing factors may include birth of a large baby, a prolonged labor, or a history of many births. A full bladder can also cause uterine atony.

Explain to the participants that a hard uterus feels like the forehead and soft uterus feels like the tip of the nose. When performing uterine massage, explain what you are doing to the mother and why. It may be uncomfortable for the mother. Use a cupped palm and massage the uterus in a circular motion until it feels contracted.

Ask the participants which uterotonics they have available for use in their facility. Discuss with them any issues they may have regarding use: stock-outs, storage, comfort with use, and/or acceptance by women, for example.

Give oxytocin

Available in
Kits 6 & 11

Oxytocin as 10 IU/mL in 1 mL ampoule

Initial dose	Continuing dose	Maximum dose
10 international units (IU) IM or IV	repeat 10 IU IM or IV after 20 minutes if heavy bleeding persists	
20 international units (IU) in 1 liter IV infusion at 60 drops/min	10 IU in 1 liter IV infusion at 30 drops/min	Not more than 3 liters of IV fluids with oxytocin

Pregnancy, childbirth, postpartum and newborn care: a guide for essential practice – 3rd ed.

Oxytocin is the medicine of choice used to help uterine atony. It causes the uterus to contract.

It can be given in the muscle of the deltoid or anterior thigh after the birth of the baby, if actively managing the third stage of labor or after the birth of the placenta. Oxytocin needs to be kept at 25°C or less. It is shipped via cold chain with Inter-Agency RH Kits 6 and 11.

Put active management of the third stage of labor (AMTSL) in the parking lot because you will have a broader discussion about the topic later in the training.

Give misoprostol

Available in
Kit 8

- Used where oxytocin is not available
- Also causes the uterus to contract
- Tablets – 200 micrograms (mcg)
- Dose = 800 micrograms (mcg) under the tongue (sublingual)

Pregnancy, childbirth, postpartum and newborn care: a guide for essential practice – 3rd ed.

Oxytocin remains the drug of choice for prevention of postpartum hemorrhage and should be provided when available. Misoprostol can be used where oxytocin is not available, or where the provider is not able to give injections. It is available in Inter-Agency RH Kit 8.

Misoprostol is a medication that also causes the uterus to contract and does not need to be kept at less than 25°C. It is more stable in hot climates.

The correct dose of misoprostol is 800 mcg sublingually (under the tongue). It comes in 200 mcg tablets. Give 4 tablets. Some sources will also say that it can be given rectally. The use of misoprostol for the prevention of postpartum hemorrhage

has been shown to be as effective as oxytocin, when given by a trained provider. Side effects are more common with misoprostol. The mother should be told what to expect: shivering, nausea, diarrhea, and fever.

Untrained providers should give misoprostol **only after** delivery of the placenta.

Ergometrine


Available in
Kit 11

- IM or IV 0.2 mg ergometrine injection
 - Max 5 doses at 15 minute intervals
- Use only after placenta is out
- Use only if oxytocin does not stop bleeding
- Associated with increased blood pressure, nausea, and vomiting
- **Do not use** in the case of eclampsia, pre-eclampsia, hypertension, or retained placenta

Pregnancy, childbirth, postpartum and newborn care: a guide for essential practice – 3rd ed.


Ergometrine is included in Inter-Agency RH Kit 11. It is **not** recommended as the first line medication for treatment of postpartum hemorrhage. Unlike oxytocin, it is associated with increased blood pressure, nausea, and vomiting. Do not use this medication to treat a patient with eclampsia, pre-eclampsia, hypertension, or a placenta that has not yet been delivered. Storage of ergometrine can also be an issue. The medication deteriorates faster than oxytocin in tropical conditions.

Bimanual compression



If heavy postpartum bleeding persists despite uterine massage, uterotonics, and removal of the placenta, then perform bimanual compression. To perform, wear sterile or clean gloves. Introduce the right hand into the vagina and clench into a fist, with the back of the hand directed posteriorly and the knuckles in the anterior fornix. Place the left hand on the abdomen behind the uterus and squeeze the uterus firmly between the two hands. Continue compression for five minutes or until the bleeding stops. If bleeding persists when bimanual compression is removed after a maximum of five minutes, apply aortic compression and transport the woman to the hospital.

Aortic compression



Feel for the femoral pulse, then apply enough pressure above the umbilicus with a clenched fist to stop the bleeding. Apply sufficient pressure until the femoral pulse is not felt. After finding the correct site, show an assistant or relative how to apply pressure. Continue pressure throughout transport to the hospital.

PPH Prevention

- **Active management of the third stage of labor (AMTSL) prevents PPH**
- **Includes:**
 1. Administration of uterotonic as soon as the baby is born
 2. Delivery of placenta via controlled cord traction
 3. Uterine massage after the placenta is delivered



It is important to carefully monitor the mother for bleeding, especially in the first two hours after childbirth.

It may not be possible to offer the full package of interventions for AMTSL in emergencies because of a lack of skilled staff, difficulty in ensuring safe injection practices, and/or a lack of refrigeration.

1. Administration of a uterotonic is the most important component of AMTSL.
2. Delivery of the placenta by controlled cord traction is recommended where skilled birth attendants are available and the provider and mother agree that a small amount of blood loss is important.
3. Postpartum abdominal uterine tone assessment is recommended for all women. Perform the uterine massage after the placenta is delivered until firm; sustained massage is not necessary.
4. Early cord clamping is no longer a component of AMTSL.

Ask participants: How do you practice in your work setting? Are there any changes that can be made to the care you provide based on the resources available? What is needed to provide AMTSL for all women?

Unit 4: Manual Removal of the Placenta

Time

60 minutes

Objectives

By the end of this unit, participants will be able to:

- ◆ Demonstrate competence in active management of the third stage of labor (AMTSL).
- ◆ Recognize indications for manual removal of the placenta at multiple levels of care.
- ◆ Explain the procedure for manual removal of the placenta.

Materials

- ◆ Video - manual removal of placenta (available on the USB key and IAWG website)
- ◆ *Active Management of Third Stage of Labor Skills Checklist* and *Manual Removal of Placenta Skills Checklist* handouts, one each per participant
- ◆ Presentation
- ◆ Normal delivery supplies (see Course Materials Checklist)

Instructions

Start this unit by reviewing the normal physiology of AMTSL and signs of placental separation using the **presentation** slides. After reviewing information on heavy bleeding before the birth of the placenta, prepare to **show the video** on manual removal of the placenta. The video is 15 minutes long and reviews the indications for and the procedure of manually removing the placenta.

After the video, **ask the participants:**

- ◆ Do you have experience with removing the placenta manually?
- ◆ Are the resources available at your respective facilities for managing this emergency?
- ◆ If not, then where would the patient be transported?

Briefly review the slides on indications for manual removal of the placenta and follow-up care after manual removal.

Emphasize that manual removal of the placenta should take place in a referral hospital and can only be performed outside the hospital to save the mother's life in an emergency.

Move on from the presentation about the prevention of postpartum hemorrhage and retained placenta to discuss AMTSL. AMTSL is a central component of reducing postpartum hemorrhage. It has been shown to reduce postpartum hemorrhage in over 60% of women and decreases the risk of retained placenta.

After reviewing the steps of AMTSL, **ask participants:**

- ◆ How do you practice in your work settings?
- ◆ Are there any changes that can be made to the care you provide based on the resources available?
- ◆ What is needed to provide AMTSL for all women?

Exercise

Demonstrate the steps of active management using the childbirth simulator or similar. Make sure to clearly demonstrate each step listed on the *Active Management of Third Stage of Labor Skills Checklist* (available on the USB key and IAWG website).

ACTIVE MANAGEMENT OF THE THIRD STAGE OF LABOR SKILLS CHECKLIST	YES	NO
1. Following delivery of the infant, the provider checks for a second baby		
2. Tells the woman what medication she is being given		
3. Gives uterotonic medication within 1 minute of delivery of the infant		
4. Tells the patient why she is getting the medication		
5. Cuts the cord. Observe for (1) changing/or taking off the first pair of gloves and (2) cutting after giving the uterotonic.		
6. Applies counter-pressure while performing controlled cord traction		
7. Only performs controlled cord traction when the patient is having a contraction		
8. Uses both hands to catch the placenta		
9. Gently turns the placenta while it is being delivered		
10. Assesses fundal tone immediately following the delivery of the placenta		
11. Inspects the placenta and membranes for completeness		
12. Checks the woman's bleeding		

Adapted with permission from Jhpiego. Helping Mothers Survive: Bleeding after Birth. 2015.

Presentation



Unit 4: Manual removal of the placenta

Basic emergency obstetric and newborn care
in humanitarian settings



Signs of placenta separation

- Third stage of labor
- Normal gush of blood before the placenta
- Lengthening of the cord
- Change in position of the uterus
- If no signs after 30 minutes:
 - Maintain skin-to-skin, breastfeeding
 - Empty bladder
- If no signs after 60 minutes:
 - Transfer to a higher level of care



It is normal to see a gush of blood from the vagina when the placenta separates from the uterine wall. The cord will also look longer as it drops closer to the vaginal opening. The top of the uterus will rise to the umbilicus or above. Gently push the uterus upward from just above the pubic bone to see if the placenta is still attached. If the cord moves back up into the vagina, the placenta may not have separated.

The mother and baby will already be skin-to-skin. Increased nipple stimulation from the baby via breastfeeding or hand motions will help the mother to release oxytocin and facilitate the delivery of the placenta.

Perform controlled cord traction to deliver placenta

- Must be gentle and only during a contraction
- Always stabilize the uterus
- Only pull the cord in a downward, steady direction



- If the uterus moves downward
- If the cord is tearing
- If the pulling hurts the mother

After there have been signs of placental separation, wait for a contraction and provide counter-traction to stabilize the uterus by using one hand to guard the uterus just above the pubic bone. Pull gently down on the cord with the other hand during a contraction with a downward, outward motion while the mother pushes.

As the placenta comes out, twist it to keep the membranes from tearing and remove slowly from the vagina. When the placenta is delivered, check to make sure the placenta and accompanying membranes are complete. Retained fragments can cause the uterus to not contract.

Stop pulling on the cord if you notice signs that the placenta has not separated. Pulling on the cord when the placenta is still attached or when the uterus is not contracted is dangerous. The cord can break (evulse) or the uterus can come out of the body, both of which can be life-threatening emergencies.

If bleeding heavily before placenta has separated

- Breastfeed, empty bladder
- Feel abdomen for a second baby
- Oxytocin 10 units in side of thigh muscle **OR** 600 mcg misoprostol by mouth
- Gentle controlled cord traction with contraction
- Prepare for transport and to treat for shock



Constant bleeding before the placenta has separated is not normal. If the mother is bleeding heavily after the birth of the baby and before the placenta has separated, it is important to try to deliver the placenta quickly. If out of the hospital, this emergency is an indication to deliver the placenta manually. Note that the dose of misoprostol is 600 mcg by mouth (oral administration) before delivery of the placenta, as opposed to 800 mcg sublingually for postpartum hemorrhage.

Pause the presentation here and **play the accompanying video** on manual removal of the placenta (available on the IAWG website and USB key).

After the video, **ask the participants** if they have experience with removing the placenta manually. Are resources available at their respective facilities for managing this emergency? If not, then where would the patient be transported?

Manual removal of the placenta

- Only in an emergency to save a mother's life
- Transfer to a higher level of care if not bleeding heavily
- Give antibiotics if transport to hospital will take more than 1 hour
 - Amoxicillin 1g by mouth, once (250 mg x 4)
 - Metronidazole 1 g by mouth, once (250 mg x 4)



Remove the placenta manually if you think the woman will bleed to death before you can reach a higher level of care. For example, if heavy vaginal bleeding continues after the massage and oxytocin and the placenta cannot be delivered by controlled cord traction, or if the placenta is incomplete and bleeding continues after administering oxytocics and applying compression.

If the placenta needs to be removed manually and the mother is **not bleeding heavily**, then plan to transfer the mother to a higher level of care as soon as possible. At the hospital, the mother will be able to receive appropriate pain medication, antibiotic therapy, and monitoring after the procedure.

Give medicine to prevent infection if it will take more than one hour to get to the hospital or other advanced help. Oral tablets are available in Inter-Agency RH Kit 6.

After manual removal

- Repeat 10 units oxytocin IM or IV
- Massage the uterine fundus
- Give ampicillin 2g IM or IV
- Give fluids slowly for 1 hour after removal (if bleeding stops)
- If bleeding continues → **urgent referral**, 20 units oxytocin per liter of IV fluids rapidly



After this slide, begin the demonstration of AMTSL and have the participants follow along using the checklist.

Unit 4 Activities

Time

1 hour and 30 minutes

Objectives

By the end of this unit, participants should be able to:

- ◆ Demonstrate competence in AMTSL.
- ◆ Demonstrate an understanding of the steps of manual removal of the placenta.

Divide the participants into two groups. Individuals in the first group will take turns demonstrating the steps of AMTSL while the performance is marked off on the *Active Management of Third Stage of Labor Skills Checklist*. Each participant will have four minutes and feedback will be given to the entire group during the debriefing. The second group will take turns demonstrating the steps for manual removal of the placenta while the performance is marked off on the *Manual Removal of the Placenta Skills Checklist* (both checklists are available on the USB key and IAWG website).

MANUAL REMOVAL OF PLACENTA SKILLS CHECKLIST		YES	NO
Preparation			
1	Explain the procedure to the woman and provide emotional support		
2	Insert an IV line with fluids rapidly infusing (active bleeding)		
3	Assist the woman onto her back		
4	Give diazepam * 10 mg IM/IV		
5	Clean vulva and perineal area		
6	Ensure bladder is empty; catheter if necessary		
7	Wash hands and forearm well and put on sterile gloves (long if available)		
Technique			
8	With one hand, hold the umbilical cord at the clamp; pull cord gently until horizontal		
9	Insert other hand into the vagina and follow the cord up into the uterus		
10	Drop the cord with external hand, hold the fundus, provide counter-traction, and prevent inversion		
11	Move fingers of internal hand sideways to locate the edge of the placenta		
12	Keep fingers tightly together and use edge of hand to gradually make space between the placenta and uterine wall		
13	Proceed gradually until the entire placenta is detached from the uterine wall		
14	Withdraw internal hand from the uterus gradually, bringing the placenta with it		
15	Use the internal hand to explore the inside of the uterine cavity to ensure all placental tissue has been removed		
16	Examine the uterine surface of the placenta to ensure lobes and membranes are complete		
17	Give prophylactic antibiotics *Ampicillin 2g IV or IM		

Adapted from WHO Pregnancy, Newborn and Childbirth Care: A guide for essential practice. 2015. 3rd Edition.

Unit 5: Transport and Referral

Time

45 minutes

Objectives

By the end of this unit, participants should be able to:

- ◆ Safely stabilize and prepare a patient for transport after postpartum hemorrhage.
- ◆ Practice IV access and fluid administration.

Materials

- ◆ Presentation
- ◆ Normal delivery supplies (see Course Materials Checklist)
- ◆ IV supplies

Instructions

This unit addresses how to safely transport and refer a patient to a higher level of care during or after a postpartum hemorrhage. The ability to safely transfer care of a patient experiencing an obstetric emergency needs to be addressed for each of the conditions discussed in this module. This unit serves as a thorough review of the process. It will subsequently be returned to for discussion with other conditions as the module progresses. Explain that the principles of transport and referral will apply to a patient being transported for any obstetric emergency.

Start this unit with a **discussion about patient transport**:

- ◆ Where do patients go if they have an obstetric emergency?
- ◆ Who arranges transportation?
- ◆ Is there a protocol in place at the facility for transferring a patient?
- ◆ If not, who would be responsible for setting up this system?

Move through the slides, addressing the principles of effective referral. The dialogue will then shift to IV fluid administration, which is crucial for safely transporting patients with active bleeding. **Demonstrate** how to appropriately start an IV by walking through the steps of the *IV Skills Checklist*. **Ask the participants** how frequently they start IVs to gauge their experience with this skill. There is time built into the end of the day to act as skills building time for IV administration. **Complete a skills checklist** for each participant if the participants state that they do not feel confident with IV administration.

INTRAVENOUS (IV) SKILLS CHECKLIST		YES	NO
1	Explain the procedure to the woman and provide emotional support		
2	Gather all the supplies <ul style="list-style-type: none"> ❖ sterile IV fluids ❖ sterile plastic tubing ❖ sterile IV ❖ tape ❖ skin antiseptic 		
3	Wash hands with soap and water; wear clean gloves		
4	Open the package of sterile tubing and attach it to the fluids using sterile technique		
5	Hang up the bag of solution high enough to let it flow to the end of the tubing		
6	Tie the tube off at the end to prevent fluid waste		
7	Use cloth or a rubber tourniquet on the upper arm		
8	Identify a vein and clean the skin		
9	Hold the vein steady between the first finger and thumb with one hand, and carefully insert the needle into the vein with the other hand. Look for blood in the needle hub. Lay the needle almost flat and slide it the rest of the way into the vein.		
10	Remove the tourniquet and attach a fluid tube to the needle		
11	Use tape to hold the needle in place		
12	Regulate the amount and speed of IV fluid flow depending on the medical condition		

Presentation



Unit 5: Transport and referral

Basic emergency obstetric and newborn care
in humanitarian settings



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISIS

This unit provides a review of transport and referral for a patient who needs to move to a higher level of care during or after postpartum hemorrhage. Explain that the principles of transport and referral will apply to a patient being transported for any obstetric emergency.

Discuss: Where do patients go if they have an obstetric emergency? Who arranges transportation? Is there a protocol in place at the facility for transferring a patient? If not, who would be responsible for setting up this system?

Referral

- **Organize reliable transportation**
- **Communicate with the receiving facility**
 - Explain her diagnosis and condition
 - Describe care provided, including medication
 - Estimate her time of arrival
- **Accompany by a provider and companion**
 - Monitor fetus/newborn
 - Fundal massage
 - Maintain IV fluids



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISIS

Discuss: What are the steps in executing an effective referral? Guide learners to the following responses:

- ◆ Organize reliable transportation.
- ◆ Communicate with the referral facility:
 - ❖ Explain her diagnosis and condition.
 - ❖ Describe care provided, including medicines.
 - ❖ Estimate her time of arrival.
- ◆ Ensure that the woman is accompanied by a companion.
- ◆ If possible, send a provider to go with her to:
 - ❖ Monitor her and the fetus/newborn.
 - ❖ Check for uterine atony.
 - ❖ Keep her IV in place with fluids infusing as required. Fluids are infused rapidly if active, heavy bleeding. Fluids are infused slowly if the bleeding is slow or the patient is stabilized.
 - ❖ Complete the referral record to send with her.
- ◆ Record the referral.
- ◆ Obtain the report from the referral facility and ensure that she receives follow-up care.

Transfer

Patient Transfer Record

Date: _____

Patient name: _____ Date of Birth: _____

Reason for transfer: _____

Date and time of delivery: _____

Medications given and time: _____

Time transfer initiated: _____ Vital signs at transfer: _____

Contact information of transferring provider: _____

Provider name: _____ Signature: _____

It is best practice to maintain documentation of a patient transfer for appropriate follow-up and organizational review. This slide has an example of a patient transfer record.

IV fluids

- Use a 16 or 18 g needle
- Run normal saline or Ringer's lactate solution
- Rapidly infuse fluids if in shock
 - 1 liter as fast as possible (15-20 minutes)
 - 1 liter at 30 mL/min (30 min)
 - Monitor BP and pulse every 15 min
 - Watch for shortness of breath or swelling
 - Reduce to 3mL/min (6-8 hours) when pulse is <100 bpm or systolic BP >100 mmHg



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISIS

World Health Organization. 2015. Pregnancy, childbirth, postpartum and newborn care: a guide for essential practice – 3rd ed.

Give IV fluids at a **rapid rate** if the patient in shock, systolic blood pressure (BP) is <90 mmHg, pulse >110/minute, or she has heavy vaginal bleeding.

Reduce the infusion rate to 3 mL/minute (1 liter in 6-8 hours) when pulse slows to less than 100/minute, or systolic blood pressure increases to 100 mmHg or higher.

Reduce the infusion rate to 0.5 mL/minute if breathing difficulty or puffiness develops.

It is important to monitor urine output and the total amount of IV fluids given. Record the time and amount of fluids given during the transfer.

IV fluids

Slowly infuse fluids if pre-eclampsia/eclampsia

- Infuse 1 liter in 6-8 hours (3 mL/min)

Moderately infuse fluids if dehydration/fever or severe pain

- Infuse 1 liter in 2-3 hours

Oral rehydration solution if no IV available

- 300-500 mL/hour



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISIS

World Health Organization. 2015. Pregnancy, childbirth, postpartum and newborn care: a guide for essential practice – 3rd ed.

- ◆ Give fluids at a **slow rate** if severe anemia/severe pre-eclampsia or eclampsia are present. It is not safe to rapidly infuse fluids into a woman with elevated blood pressure.
- ◆ Give fluids at a **moderate rate** if severe abdominal pain, dangerous fever, or dehydration is present.
- ◆ Give oral rehydration solution (ORS) by mouth if the woman is able to drink. Never give ORS to a woman who is unconscious or having convulsions.

Unit 6: Antibiotics for Postpartum Infection

Time

1 hour and 45 minutes

Objectives

By the end of this unit, participants will be able to:

- ◆ Review and apply assessment, diagnosis, treatment, and evaluation of postpartum infection.
- ◆ Appropriately identify and refer women to a higher level of care for severe infection.

Materials

- ◆ *Case Study – Fever after childbirth* handout, one per participant
- ◆ Normal delivery supplies (see Course Materials Checklist)
- ◆ Simulated medications – antibiotics
- ◆ IV supplies: Infusion sets, IV cannulas, bags of IV fluids

Instructions

Open this unit with a **role play**. As a postpartum woman, come in acting very ill and complain of pain and fever. Act minimally responsive and weak. Have the co-facilitator state that an IV is needed.

Ask the learners:

- ◆ Have you ever seen a woman come in after having a baby who is very sick?
- ◆ What happened?
- ◆ Is there any other treatment that could have been available if the woman had delivered in a hospital? Is there access to IV supplies at your health facilities?

Wrap up the discussion and start the slide **presentation**. Go through the slides and discuss the related notes. Finish the slides related to treatment at the referral level of care. Explain that later in the afternoon, there will be an opportunity to practice with IV infusion supplies.

Case Study

Utilize the prompt and the following instructions for the case study activity:

- ◆ Divide participants into small groups of two to three and ask the learners to go through the case study together.
- ◆ Give the groups 30 minutes to go over the case study and then come back together as a group to discuss the case.

Take a short break after debriefing from the case study before starting the next unit.

Case Study: Fever After Childbirth

Mrs. E is a 35-year-old para three. Mrs. E's husband has brought her to the health center today because she has had a fever and chills for the past 24 hours. She gave birth to a full-term infant at home 72 hours ago. Her birth attendant was the local traditional birth attendant (TBA). Labor lasted two days and the TBA inserted herbs into Mrs. E's vagina to help speed up the childbirth. The newborn breathed spontaneously and appears healthy.

Assessment (history, physical examination, screening procedures/laboratory tests)

1. What will you include in your initial assessment of Mrs. E? Why?
 - ❖ History of labor
 - ❖ Presence of chills/fever/pain/vaginal bleeding
 - ❖ History of malaria, HIV, or anemia
 - ❖ Vital signs
 - ❖ Uterine tone and tenderness
2. What particular aspects of Mrs. E's physical examination will help you make a diagnosis or identify her problems/needs? Why?
 - ❖ Fever greater than 38°C
 - ❖ Chills or malaise
 - ❖ Uterine tenderness or subinvolution
 - ❖ Purulent or foul lochia
 - ❖ Signs of shock
 - ❖ Pulmonary edema

Diagnosis (identification of problems/needs)

You have completed your assessment of Mrs. E and your main findings include the following:

History: Mrs. E admits that she has felt weak and lethargic, has abdominal pain, and has noticed a foul-smelling vaginal discharge. She does not have pain when she urinates and does not live in an area with malaria. She had a tetanus immunization and a booster 3 years ago. It is unknown whether her placenta was complete.

Physical Examination

- ❖ Vital signs: temperature 39.8°C, pulse 136 beats per minute, respirations 24 per minute
 - ❖ Pale, lethargic, and slightly confused
 - ❖ Uterus at 1 cm below the umbilicus and tender
 - ❖ Foul-smelling vaginal discharge, no tears or lesions
 - ❖ Cervix is 2 cm dilated with cervical motion tenderness
3. Based on these findings, what is Mrs. E's diagnosis (problem/need)? Why?
 - ❖ Puerperal sepsis due to elevated temperature and abdominal tenderness. The lethargy and confusion raise suspicion for sepsis.

Care provision (planning and intervention)

4. Based on your diagnosis (problem/need identification), what is your plan of care for Mrs. E? Why?
 - ❖ Draw/send blood cultures if lab facilities are available
 - ❖ Start IV fluids, first dose antibiotics, and prepare to transport/refer
 - ❖ Plan on IV antibiotics x 48 hours fever free
5. Based on these findings, what is your continuing plan of care for Mrs. E? Why?
 - ❖ Antipyretics while febrile
 - ❖ Uterine evacuation, if needed
 - ❖ Bed rest, perineal hygiene, newborn with mother

Presentation



Unit 6: Antibiotics for postpartum infections

Basic emergency obstetric and newborn care
in humanitarian settings



Open the unit with a **role play**. Follow the instructions in the facilitator's guide for more information.

Clean delivery

- Standard precautions in health facilities
- Clean birth practices in health facilities
- Clean delivery kit distribution if delivery outside health facility



In humanitarian settings, there may be a lack of health supplies or infrastructure, and an increased workload. Staff working in the health sector may resort to taking shortcuts in procedures, which endanger the safety of both patients and staff. Therefore, it is essential that standard precautions are respected. **Ask the participants:** How do we reduce the risk of infection during and after childbirth? What are standard precautions? Answers include:

- ◆ Frequent hand-washing
- ◆ Wearing appropriate gloves – sterile single-use gloves for internal examinations, non-sterile single use gloves for handling contaminated materials
- ◆ Wearing protective clothing
- ◆ Safe handling of sharp objects
- ◆ Proper waste disposal
- ◆ Instrument sterilization
- ◆ Housekeeping

Distribution of clean delivery kits to visibly pregnant women in the population gives women who are delivering without the presence of a skilled birth attendant the tools needed to reduce the risk of infection.


History

- Always **take the history** of the patient first
- Long labor? Premature rupture of membranes (PROM) >18 hours? Manual removal of placenta? C-section? PPH? Traditional practices?
- Malaria? HIV? Anemia?
- Chills? Fever? Pain? Bleeding?



When a woman presents with a possible infection, it is important to always take the history of the patient first. Discuss her history to gain an initial clinical picture of the situation. Check any antenatal record or delivery record available. Some women are more vulnerable to puerperal sepsis, including those who are anemic and/or malnourished.

Protracted labor, prolonged rupture of membranes, frequent vaginal examinations, a traumatic delivery, cesarean section, and retained placental fragments all predispose to puerperal infection.



The physical exam will give you further information about the status of the patient.

Vital signs and general condition: Does the patient have signs of shock, including a rapid pulse? Is there a fever greater than 38 C? Is the patient alert and oriented? Is she shaking with chills and fatigued? Is her breathing easy or labored? Do her lungs sound clear and are her respirations normal?

Abdominal tenderness: Is her abdomen tender when you palpate for her uterus? Is the uterus palpable in her abdomen? Where do you find it? Is it larger than you would expect it to be? Is the uterus hard or soft? When did she last empty her bladder?

Pelvic exam: A gentle internal exam will give more information on uterine size and tenderness, the presence of tears, lochia odor, and the amount of lochia.

Breasts: Are her breasts engorged with redness or other signs of an infection?

Differential diagnoses

- Urinary tract infection (bladder or kidneys)
- Wound infection
- Mastitis or breast abscess
- Thromboembolic disorder
- Pneumonia
- Malaria or typhoid
- HIV-related infections



Based on the assessment of her obstetric history and targeted exam, make your diagnosis. If the patient does not fit the clinical picture described on the previous slide, then move through the other differential diagnoses.

Treatment

- **Stabilize and refer**
- **Give first dose of antibiotic prior to referral**
- **Antibiotic therapy until 48 hours fever free**
 - Clindamycin 150 mg every 6-8 hours
 - Gentamicin 80 mg IM every 8 hours
- **Send blood cultures, if lab facilities are available**



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISIS

Oral antibiotics are not effective for treating a severe postpartum infection. The mother will need IV antibiotic therapy at a facility with appropriate monitoring capacity. Outside of the hospital, a patient presenting with a severe infection will need emergency treatment with a first dose and immediate referral to a higher level of care.

The mother needs broad spectrum antibiotics until 48 hours fever-free. The Inter-Agency RH Kits for the health facility level only contain oral antibiotics. The Kits are in the process of being updated to reflect current guidelines. **Ask the participants:** which antibiotics are available at your facility?

Lab facilities will not be available most of the time.

Treatment

- **Medicine to reduce temperature**
- **Uterine evacuation if needed (MVA)**
- **Consider possible tetanus (herbs in vagina?)**
- **Bed rest and perineal hygiene**
- **Keep newborn with mother – monitor closely**



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISIS

A patient may need further treatment upon arrival at the hospital. Ongoing treatment may include the following, as listed on this slide:

- ◆ Paracetamol 500mg tablets can be given every 4 to 6 hours for pain and fever management.
- ◆ Uterine evacuation using manual vacuum aspiration (MVA) can be performed if there are retained products from the pregnancy contributing to the illness.
- ◆ Tetanus toxoid may be indicated if the patient has had herbs inserted into the vagina during labor and birth, and has not completed prior immunization.

Referral

- **Organize reliable transportation**
- **Communicate with the receiving facility**
 - Explain her diagnosis and condition
 - Describe care provided, including medication
 - Estimate her time of arrival
 - Send a referral letter with her
- **Accompany by a provider and companion**
 - Monitor newborn
 - Ensure the first dose of antibiotics is given
 - Maintain IV fluids



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISIS

Discuss: What are the steps in executing an effective referral? Guide learners through the following responses:

- ◆ Organize reliable transportation.
- ◆ Communicate with the referral facility:
 - ❖ Explain her diagnosis and condition.
 - ❖ Describe care provided, including medicines.
 - ❖ Estimate her time of arrival.
- ◆ Ensure that the woman is accompanied by a companion.
- ◆ If possible, send a provider to go with her to:
 - ❖ Monitor her and the fetus/newborn.
 - ❖ Keep her IV in place.
 - ❖ Complete the referral record to send with her.
- ◆ Record the referral.
- ◆ Obtain a report from the referral facility and ensure that she receives follow-up care.

Unit 7: Magnesium Sulfate for Severe Pre-eclampsia/Eclampsia

Selected components of this unit have been adapted with permission from Jhpiego. Helping Mothers Survive – Pre-eclampsia and eclampsia. 2016.

Time

2 hours and 30 minutes

Objectives

By the end of this unit, participants will be able to:

- ◆ Understand the continuum of hypertensive disorders in pregnancy.
- ◆ Demonstrate an ability to accurately measure and record blood pressure.
- ◆ Demonstrate an ability to assess for severe pre-eclampsia and eclampsia in limited resource settings.
- ◆ Demonstrate an ability to safely prepare magnesium sulfate for intramuscular (IM) and IV administration.
- ◆ Understand treatment protocols for antihypertensive medication administration.

Materials

- ◆ Normal delivery supplies (see Course Materials Checklist)
- ◆ Simulated medications – magnesium sulfate, 2% lignocaine, calcium gluconate, and hydralazine
- ◆ Needles for IM injection
- ◆ Bladder catheter
- ◆ Reflex hammer
- ◆ Urine dipsticks
- ◆ Melon or other local fruit
- ◆ Sharps container

Instructions

Open this unit with a **role play** of a one-day postpartum woman complaining of a severe headache who begins to convulse and loses consciousness. If you are facilitating alone (not recommended), ask a learner in advance to play the woman and ask for help from the participants, but do not interfere.

After the role play, guide the **discussion**:

- ◆ Has anyone seen an eclamptic convulsion or a woman die from eclampsia?
- ◆ What happened?
- ◆ Was there anything that could have been done to help?

Note to Trainers: This refresher training assumes a baseline level of familiarity with management of pre-eclampsia and eclampsia. If the need for further training on these issues is identified, then give appropriate feedback to the training coordinator.

Start the **presentation** by reviewing the slides that classify the hypertensive disorders of pregnancy. Explain that this module will review the management of severe pre-eclampsia and eclampsia using magnesium sulfate and anti-hypertensives. Pause after the slide about accurate blood pressure measurement and complete the blood pressure activity below, if appropriate for the participants in the group.

Optional: Blood Pressure Measurement Activity

Ask the participants to work in pairs to practice blood pressure measurement. Make sure you have a stethoscope and blood pressure machine for each pair. Pay attention to the learners' techniques. Provide supportive correction as needed. Once the first participant has taken a reading, ask each pair to share their measurements. Note if 120/80 or 130/80 are overrepresented. If so, ask the learners if they note anything unusual about the blood pressures. Encourage them to become "expert" blood pressure takers and then have them switch roles.

Review the slides on the classification of pre-eclampsia, severe pre-eclampsia, and eclampsia and the treatment of severe pre-eclampsia/eclampsia. On the role play slide that includes the classification of the disorders, **explain the activity** listed below to the participants.

Activity

Adapted with permission from Jhpiego. Helping Mothers Survive – Pre-eclampsia and eclampsia. 2016.

Instructions

Divide the group into pairs. Ask one learner to play the role of the woman. Ask the other learner to play the role of the provider. Have the participants come to one side of the room to receive instructions privately. Ask them to bring paper and pencil to record their measurements.

Give the participants this information privately:

- ❖ 25 year-old gravida 1 para 0 (G1P0) at 37 weeks gestation
- ❖ Blood pressure is 160/102
- ❖ Urine 2+ protein
- ❖ Danger signs: no headache, no visual changes, no right upper quadrant pain, no difficulty breathing, and no rales

Tell them to only provide this information in response to measurements taken by providers. Tell providers they must take the actual measurements during the role play, not simply talk about what they would do. Send the participants back to their stations where they will present for care.

Read the following paragraph before they begin:

Mrs. M is presenting to you in very early labor. Please interview her and proceed normally for a routine labor evaluation. If you are taking her blood pressure, listening to her lungs, or measuring her urine for protein, then do so. Your client will tell you the measurement.

Observe the role plays. Ensure that participants stay on track, but do not correct or interrupt the role play.

When all the teams are done with the assessment, debrief with the participants. Ask: What is your most likely diagnosis? Why?

The correct answer is **severe pre-eclampsia** because the blood pressure is in the severe category and she has 2+ protein.

Demonstration and Activity: Preparation and Administration of Magnesium Sulfate

Set up the supplies for the activity on preparation and administration of magnesium sulfate during the break, including the following:

- ☐ Magnesium sulfate 500 mL/10 mL vials, at least 3 vials
- ☐ Lignocaine 2%
- ☐ Sterile water
- ☐ 20 cc syringes, at least 3
- ☐ IM needles, at least 3
- ☐ IV normal saline (NS) or lactated ringers (LR)
- ☐ IV infusion set and IV start kit (IV needle, alcohol swabs, gloves, tourniquet, tape)
- ☐ Calcium gluconate 10%
- ☐ Hollowed out melon or other local fruit with skin for IM injection
- ☐ Sharps container

Bring the group back together and start the slide **presentation** with magnesium sulfate. It is important to emphasize the safety concerns around giving magnesium sulfate outside of a facility with appropriate monitoring. The slide on the formulation of magnesium sulfate available in the Inter-Agency RH Kits discusses the difference between IM and IV solution concentrations. **Magnesium sulfate must be diluted with sterile water to a 20% solution prior to administering intravenously.**

Demonstrate how to draw up the loading dose of magnesium sulfate for IM injection by appropriately drawing 5 g in 10 mL of the 50% solution and the 1 mL of lignocaine in the same syringe. Demonstrate where the injection is given in the upper outer buttock.

Ask a volunteer or have the co-facilitator set up the IV infusion set.

Demonstrate how to first dilute the magnesium sulfate with sterile water prior to IV injection.

Review the IV administration of a repeat dose of magnesium sulfate and the indications to give or hold the subsequent dose. **Discuss** with the participants how they safely monitor IV infusions in their facility. Are they able to do slow IV injections? If not, then where will they refer a patient after the IM loading dose? Briefly discuss the recommended use of diazepam as a secondary medication if magnesium sulfate is not available or convulsions recur after a second dose of IV medication.

Calcium gluconate is the antidote to magnesium sulfate. If the patient suffers respiratory arrest during the administration of magnesium sulfate, stop the infusion immediately and then give 1 gram of calcium gluconate IV over 10 minutes.

Ask the participants:

- ◆ Where are the magnesium sulfate and calcium gluconate stocked in your facility?
- ◆ How are they monitored?

Activity: Preparing and administering magnesium sulfate (MgSO₄)

Invite the participants to the table and have them work in groups to practice using magnesium sulfate IM and IV, and to draw up calcium gluconate. Use the melon or other local fruit to practice deep IM injection.

Bring the group back together after the magnesium sulfate activity and **discuss** antihypertensive medications. Hydralazine 5 mg IV over 1 to 2 minutes is the first-line recommendation. It is currently available in Inter-Agency RH Kit 11. Discuss with the participants to understand which antihypertensives are available in their setting. Other medications that may be available include:

- ◆ **Nifedipine immediate-release only** – 5-10 mg by mouth or bitten then swallowed. Repeat every 30 minutes until the target blood pressure is achieved. Max dose 30 mg.
- ◆ **Labetalol oral** – 200 mg by mouth, repeat every one hour if needed to achieve the desired outcome. Max dose 1200 mg in 24 hours.
- ◆ **Labetalol IV** – 10 mg IV over, double dose every 10 minutes to 80 mg until goal is achieved. Max dose 300 mg, then switch to oral (do not use if patient experiences congestive heart failure, asthma, or hypovolemic shock).

Address any outstanding questions on the administration of magnesium sulfate or antihypertensives for severe pre-eclampsia and eclampsia.

Introduce the group to the clinical simulation outlined below. Tell learners that they will be participating in a simulation. Participants act as if they are providing care for a real woman. Observers will be ready to provide feedback. If there are more than three participants, then a second simulation will be needed so that all learners can actively participate.

Clinical Simulation

Adapted with permission from Jhpiego. Helping Mothers Survive – Pre-eclampsia and eclampsia. 2016.

Overview

As the facilitator, you will be Mrs. C. Assign up to 3 learners to act as providers and any remaining learners to observe and give feedback after the role play. Allow 15 minutes for this activity. The goal is for learners to diagnose and initiate treatment of Mrs. C, who has severe pre-eclampsia even though she has no danger signs and appears to be normal. Her blood pressure is dangerously high and the team needs to identify and begin treatment quickly. The providers are working in a health clinic in a rural area. Read the following scenario to the group:

Case 1: *I am Mrs. C, 27 years old and 39 weeks pregnant with my first baby. I have been to 4 antenatal care (ANC) visits with no problems and I think I am in labor. Please begin your assessment and tell me what you are doing. If you choose to assess something, please do so and I will give you the results. Afterwards we will debrief as a team.*

Give the following information, but only if and when assessed:

- ❖ Blood pressure is 178/112
- ❖ No danger signs
- ❖ Urine protein 2+
- ❖ Fetal heart rate 140 beats per minute (bpm), fetal movement present
- ❖ Sterile vaginal exam – cervix is 4 cm dilated
- ❖ Contractions every 4 minutes lasting 45 seconds, moderate
- ❖ All other measurements are normal

Debrief: Upon completion of the role play, begin a structured debrief. Be sure to include feedback about communication and respectful maternity care. Include the following points:

- ❖ Confirm correct diagnosis and treatments
- ❖ SPE and full MgSO₄ loading dose (IV and IM) and antihypertensive

- ❖ **Ask:** Why this diagnosis? What went well? Did you miss anything? How did the team function? What did you learn that you will remember for next time?
- ❖ Help learners identify that women can have SPE without danger signs. The team must move quickly from assessment and diagnosis to management.
- ❖ Be sure to address any communication gaps. Were all team members comfortable working together and delegating? As the client, share how you felt. Did they delegate, communicate, and provide respectful maternity care?

Repeat with Case 2, if needed, based on the following information:

Case 2: Read the same case but use the following measurements when assessed:

- ❖ Blood pressure is 156/106
- ❖ Blurred vision
- ❖ Urine protein 2+
- ❖ Fetal heart rate 150 bpm, fetal movement present
- ❖ Sterile vaginal exam: Cervix is 7 cm dilated
- ❖ Contractions every 3 minutes x 60 seconds, moderate
- ❖ All other measurements are normal

Bring the group together after the clinical simulation is complete and all participants have had a chance to participate. Discuss what went well and what could have been done differently.

Presentation



Unit 7: Magnesium sulfate for pre-eclampsia / eclampsia

Basic emergency obstetric and newborn care
in humanitarian settings



Open the unit with a **role play**. Follow the instructions in the facilitator's guide for more information.

Hypertensive disorders in pregnancy

- Chronic hypertension
- Gestational hypertension
- Pre-eclampsia (mild to severe)
- Superimposed pre-eclampsia
- Eclampsia



Inter-agency
RH Kit 6

Hypertension complicates 5 to 10% of all pregnancies and is associated with 14% of all maternal deaths. There is a spectrum of hypertensive disorders, which are often progressive and move quickly to severe disease. This session focuses on managing severe pre-eclampsia (SPE) and eclampsia. Severe pre-eclampsia and eclampsia are considered to be life-threatening emergencies and will require referral from a health center to a higher level of care for ongoing therapy and treatment.

Blood pressure is checked any time a pregnant/postpartum woman presents with a problem. Magnesium sulfate, the drug of choice for treatment of severe pre-eclampsia and eclampsia, is available in Inter-Agency RH Kit 6.

Assessment

- Gestational age > 20 weeks
- Accurate blood pressure reading
- Presence of protein in urine ($\geq 2+$ dipstick)
- Presence of danger signs
 - Nausea/vomiting
 - Epigastric pain
 - Headache
 - Visual changes
 - Right upper quadrant pain



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN EMERGENCIES

A diagnosis of pre-eclampsia is made after 20 weeks of gestation if blood pressure is elevated on two consecutive readings at least 4 hours apart with the woman at rest and no prior history of high blood pressure. One reading of ≥ 160 or ≥ 110 is an emergency. Blood pressure readings can be inaccurate due to many factors – technique, equipment, or stimulants.

Ask: How do you measure gestational age at your facility? Do you have pregnancy wheels? Note that at 20 weeks, the fundus can be felt at the umbilicus.

There are urine dipsticks available in Inter-Agency RH Kit 6. Check for urine protein in women with high blood pressure. Healthy pregnant women should not have more than 1+ protein in their urine. If you cannot check for protein, begin

treatment immediately for women who otherwise meet the definition of severe pre-eclampsia. Urine dipsticks may be affected by vaginal discharge, blood, or amniotic fluid. Women must be instructed to do a clean-catch to avoid contamination. Instruct the woman to clean the vulva with water, pass a small amount of urine, then place a clean, dry cup under the stream. In settings where lab facilities are available, a urine sample can be sent for a protein to creatinine ratio analysis (greater than 0.3).

All pregnant or postpartum women should be assessed for danger signs, particularly women presenting with high blood pressure. New onset high blood pressure plus any danger sign indicates severe pre-eclampsia. Diagnose and treat severe pre-eclampsia if blood pressure is slightly elevated ($\geq 140/\geq 90$) and the woman has one danger sign, as outlined on the slide. With respect to headaches, they are typically severe and not relieved by pain medication. Visual changes may appear as blurred vision or seeing lights/spots.

There may also be **clinical signs** of severe disease on assessment. These signs may include pulmonary edema – difficulty breathing and/or rales (abnormal rattling) heard when listening to lungs, low urine output (less than 400 mL in 24 hours), and abnormal lab values (low platelets, elevated serum creatinine, and elevated liver enzymes). Swelling of the legs and feet is no longer considered a diagnostic sign. There may or may not be access to laboratory facilities for monitoring urine output and doing lab tests in an emergency context.

Accurate blood pressure

- | | |
|--|--|
| <input type="checkbox"/> Feet flat on floor | <input type="checkbox"/> Slowly release the air |
| <input type="checkbox"/> Arm at level of heart | <input type="checkbox"/> sBP at first sound |
| <input type="checkbox"/> Cuff firmly on upper arm – 2 cm above elbow | <input type="checkbox"/> dBP when sound stops |
| <input type="checkbox"/> Needle at zero | <input type="checkbox"/> Do not round the number |
| <input type="checkbox"/> Stethoscope positioned | <input type="checkbox"/> Document what you hear |
| <input type="checkbox"/> Quickly inflate cuff to 180 mmHg | |



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN EMERGENCIES

Review this checklist for accurate blood pressure measurement and request a volunteer from the participants.

Demonstrate proper technique, pointing out key steps during the demonstration. Regardless of the actual blood pressure measured, state that the reading is 152/94.

Demonstrate appropriate, respectful counseling about the need to retake the blood pressure in four hours. After the demonstration, tell the volunteer the actual blood pressure.

Perform the optional blood pressure measurement activity after this slide, if using.

Classification

Disorder	Onset	Criteria
Mild to moderate Pre-eclampsia (PE)	20 + weeks	New onset high BP at 2 readings at least 4 hours apart: sBP \geq 140 mmHg OR dBP \geq 90 mmHg PLUS Proteinuria $>$ 300mg in 24 hours or 2+ on dipstick
Severe pre-eclampsia	20 + weeks	New onset high BP of sBP \geq 160 mmHg OR dBP \geq 110 mmHg with proteinuria as above OR any one of danger signs
Eclampsia	20 + weeks	PE as defined above plus convulsions or unconsciousness

**At 37+ weeks, stabilize the patient and refer for delivery.
Induction of labor is recommended at term.**

WHO recommendations for prevention and treatment of pre-eclampsia and eclampsia. WHO Press. 2011. Geneva.

All women past 20 weeks of pregnancy and all postpartum women should have their blood pressure measured when they present to the clinic for care. Pregnant women with pre-eclampsia or eclampsia will present with raised blood pressure, protein in urine, and may have other danger signs. In severe pre-eclampsia and eclampsia, the risks to the mother's health are sufficiently high to warrant delivery of the baby irrespective of its maturity.

◆ For women with mild pre-eclampsia or gestational hypertension at term, induction of labor is recommended. In women with severe pre-eclampsia at term, a policy of early delivery is recommended. Stabilize and deliver. If it is not safe to allow labor to progress for the mother and fetus in severe pre-eclampsia, then cesarean birth is recommended.

- ◆ For women with severe pre-eclampsia, a viable fetus, and between 34 and 36 (plus 6 days) weeks of gestation, a policy of expectant management may be recommended, provided that uncontrolled maternal hypertension, increasing maternal organ dysfunction, or fetal distress are absent and can be monitored.
- ◆ For women with severe pre-eclampsia, a viable fetus, and before 34 weeks of gestation, a policy of expectant management is recommended, provided that uncontrolled maternal hypertension, increasing maternal organ dysfunction, or fetal distress are absent and can be monitored.
- ◆ Induction of labor is recommended for women with severe pre-eclampsia at a gestational age when the fetus is not viable or unlikely to achieve viability within one or two weeks.

Treatment: Severe pre-eclampsia

1. **Urgent referral to hospital unless imminent delivery**
2. **If imminent delivery, then manage childbirth before referral**
3. **Magnesium sulfate to prevent convulsions – loading dose pre-departure**
4. **Antihypertensives to lower blood pressure**
5. **Teamwork and effective communication to facilitate referral**



1. A woman with a diagnosis of severe pre-eclampsia will require care and monitoring at the referral level. If your facility can provide close monitoring, getting advanced care may mean moving the woman from the clinic or outpatient department to the labor ward. If a facility cannot provide close monitoring, women with severe pre-eclampsia should be transferred as soon as possible. Always stabilize and begin appropriate treatment before transfer.
2. Magnesium sulfate is the drug of choice for convulsions. We will discuss proper administration later in the module.
3. Antihypertensives are recommended based on the diastolic blood pressure. We will also discuss proper administration of these medications later in the module.
4. Explain to the woman and her family that she needs more services and closer monitoring than your facility can provide.

Treatment: During a convulsion

- **Left lateral position, protect from fall and injury**
- **Place and secure padded tongue blades between teeth** (not during convulsion)
- **Stabilize and refer urgently to a hospital, unless delivery is imminent**
 - Keep the woman on a left side position
 - Magnesium sulfate, anti-hypertensives, repeat during transport, if needed



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISIS

Any woman who presents with convulsions is assumed to have eclampsia. It is associated with significant maternal and perinatal morbidity and mortality, and requires immediate action. Speed is important if a woman is having convulsions. The brain is deprived of oxygen during a seizure, which can permanently damage the brain. Do not leave the woman on her own. Help her into a left lateral (side-lying) position and protect her from fall and injury. Place a padded tongue blade between her teeth to prevent a tongue bite, and secure it to prevent aspiration. Never attempt to place the tongue blade during a convulsion.

Treatment: After a convulsion

- **Give available emergency drugs**
- **Start an IV infusion**
- **Stabilize and refer urgently to a hospital unless delivery is imminent**
 - Keep the woman on a left side position
 - Magnesium sulfate, anti-hypertensives, repeat during transport, if needed



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISIS

Role play

Disorder	Onset	Criteria
Mild to moderate Pre-eclampsia (PE)	20 + weeks	New onset high BP at 2 readings at least 4 hours apart: sBP \geq 140 mmHg OR dBP \geq 90 mmHg PLUS Proteinuria $>$ 300mg in 24 hours or 2+ on dipstick
Severe pre-eclampsia	20 + weeks	New onset high BP of sBP \geq 160 mmHg OR dBP \geq 110 mmHg with proteinuria as above OR any one of danger signs
Eclampsia	20 + weeks	PE as defined above plus convulsions or unconsciousness

At 37+ weeks, stabilize the patient and refer for delivery. Induction of labor is recommended at term.

WHO recommendations for prevention and treatment of pre-eclampsia and eclampsia. WHO Press. 2011. Geneva.

Divide the group into pairs. Ask one learner to play the role of the woman. Ask the other learner to play the role of the provider. Have the actors come to one side of the room to receive instructions privately. Ask them to bring paper and pencil to record their measurements. See the facilitator's guide for specific instructions.

Take a break after this role play activity is completed and you have debriefed with the group.

Magnesium sulfate

- **Drug of choice in severe pre-eclampsia and eclampsia for the prevention and treatment of convulsions**
- **Safe administration via IV and/or IM**
 - Slow IV infusion only
 - Ongoing monitoring for respiratory and other neuromuscular depression
- **Side effects**
 - Warmth during injection
 - Flushing, thirst, headache, nausea, or vomiting



Magnesium sulfate is the drug of choice in severe pre-eclampsia and eclampsia for the prevention and treatment of convulsions. It is preferred over phenytoin and diazepam, and is provided in Inter-Agency RH Kits 6 and 11. It is thought the medicine works by dilating the blood vessels in the brain – thus reducing the systolic blood pressure – and preventing or treating eclamptic seizures.

It is crucial that this medication be administered safely because of the potential for respiratory failure or death if too much is given too fast. Rapidly infusing the medicine can lead to severe consequences. Appropriate monitoring is required to ensure the safety of the patient.

It is important to counsel the patient about the side effects of the treatment.

Monitoring

- **Closely monitor for potential serious side effects**
 - Vital signs
 - Urine output – minimum 100 mL/4 hours
 - Respiratory rate – minimum 16 per min
 - Deep tendon reflexes – knee jerk present?
- **HOLD next dose until all signs are normal**



Monitor the following to keep the patient safe:

- ◆ Vital signs.
- ◆ Urine output: Decreased output is associated with magnesium toxicity. The body filters the medicine through the kidneys.
- ◆ Respiratory rate (rate should be more than 16 breaths/minute) as magnesium (Mg) depresses neuromuscular transmission.
- ◆ Deep tendon reflexes (the knee jerk should be present before the next dose is given).
- ◆ Urine output as Mg is excreted through the kidney (decreased urine output can be associated with magnesium toxicity).

Formulation

Kits 6 and 11 contain 500 mg/mL in 10 mL vials

- One vial contains 5 mg MgSO₄ = 50% solution

Route	Dose	50% Solution	20% Solution *Add 4 mL of 50% solution to 6 mL sterile water to make
IM	5 g	10 mL mag sulfate + 1 mL of 2% lignocaine	N/A
IV	4g	8 mL	+ 12 mL sterile water = 20 mL
	2g	4 mL	+ 6 mL sterile water = 10 mL



Never give 50% solution magnesium sulfate intravenously (IV)

The magnesium sulfate in Inter-Agency RH Kit 11 and Kit 6 is provided in vials with a concentration of 500 mg/mL in 10 mL (or 5 g), which is a 50% solution (i.e. 1 g in 2 mL).

It is crucial to pay attention to the strength of the medication. The magnesium sulfate can be given in the muscle from the vial in the 50% solution. It is recommended that the magnesium be mixed with 1 mL of 2% lignocaine to dull the discomfort. **The Inter-Agency RH Kits currently contain 1% lidocaine and are being revised.**

Magnesium sulfate must be diluted with sterile water before giving it through an IV.

Loading dose administration

- Start IV fluids (NS or LR) 1 liter in 6-8 hours (3 mL/min)
- Loading dose is 4 g MgSO_4 IV *plus* 10 g MgSO_4 IM
- If IV not possible, give IM only
- Use correct formulation for route of administration

Route	Dose	50% Solution	20% Solution *Add 4 mL of 50% solution to 6 mL sterile water to make
IM	5g	10 mL mag sulfate + 1 mL of 2% lignocaine	N/A One 5 g dose in each buttock = 10 g
IV	4g	8 mL	+ 12 mL sterile water = 20 mL 4 g over 20 minutes in NS or LR

Acronyms:

NS = Normal Saline

LR = Lactated Ringers

If there is no access to intravenous (IV) supplies in the facility, then the loading dose will only be given in the muscle (IM). Draw up 5 g of magnesium sulfate (MgSO_4) 50% solution and 1 milliliter (mL) of 2% lignocaine **in the same syringe**. Give one 5 g dose in the upper outer quadrant of one buttock. Repeat the procedure with a new needle and syringe in the other buttock to administer a total IM loading dose of 10 g.

If there is IV access at the facility, then 4 g of magnesium sulfate will also be given through the IV. **Remember to dilute the magnesium sulfate with sterile water prior to IV administration.** Start an IV line and slowly administer normal saline or ringer's lactate at a rate of 1000 mL in 6 to 8 hours (3 mL/min).

Continued treatment

If referral delayed, if in active labor, or if convulsions during transport:

- 5g IM of MgSO_4 every 4 hours, alternating side of administration each time
- Continue until 24 hours after delivery or last convulsion

Route	Dose	50% Solution	20% Solution *Add 4 mL of 50% solution to 6 mL sterile water to make
IM	5 g	10 mL mag sulfate + 1 mL of 2% lignocaine	N/A



DO NOT give the next dose if:

- Knee jerk reflex is absent
- Urine output less than 100 mL in 4 hours
- Respiratory rate less than 16 breaths/minute

If there is not access to IV supplies in the facility, then the loading dose will only be given in the muscle. Draw up 5 g of magnesium sulfate 50% solution and 1 mL of 2% lignocaine **in the same syringe**. Give the 5 g dose in the upper outer quadrant of one buttock. If a next dose is given four hours later, then repeat the procedure in the other buttock. Alternate sides of treatment until 24 hours after delivery or the last convulsion.

For recurrent convulsions

After 15 minutes, more IV medication can be given if fits continue

Route	Dose	50% Solution	20% Solution *Add 4 mL of 50% solution to 6 mL sterile water to make
IV	2g	4 mL	+ 6 mL sterile water = 10 mL 2 g over 20 minutes in NS or LR



Never give 50% solution magnesium sulfate intravenously (IV)

After 15 minutes, treat recurrent convulsions with a second, smaller dose of IV magnesium sulfate. Draw up 2 g/4 mL of magnesium sulfate and dilute with 6 mL of sterile water to create a 20% solution. This diluted dose can be slowly injected via IV for over 20 minutes.

Diazepam is considered to be the next step if convulsions recur after the second dose of IV magnesium. The Inter-Agency RH Kits do not have diazepam available for this purpose. Is diazepam used in your facilities?

If respiratory arrest

- Calcium gluconate is the antidote to magnesium sulfate
- Kit 6 and 11 – calcium gluconate 100 mg/mL x 10 mL (1g)
- Give IV: 1g (10 mL of 10% solution) over 10 minutes
- Assist ventilation using bag and mask



Calcium gluconate is the antidote to magnesium sulfate. It must be available whenever magnesium sulfate will be given intravenously. If the patient stops breathing during the administration of magnesium sulfate, it is crucial to act quickly and administer the medication.

Invite the participants to the table and have them work in groups of 2 or 3 to practice using magnesium sulfate IM and IV, and to draw up calcium gluconate.

Use the melon or other local fruit to practice deep IM injection.

Inter-agency RH Kit 11

Antihypertensive medications

- Given if diastolic BP is ≥ 110 mmHG
- Hydralazine 5 mg, given IM or IV (3-4 min)
- Repeat in 30 min if diastolic BP ≥ 90 mmHG
- Maximum dose 20 mg/24 hours (4 doses)



Antihypertensive drugs should be used to lower blood pressure rapidly in cases of severe pre-eclampsia and eclampsia. Use the diastolic blood pressure to determine the need for an antihypertensive.

Hydralazine is the medication of choice to lower blood pressure. It is available in Kit 11. **Ask the participants:** What types of antihypertensive drugs do you have available in your facility?

Other medications that may be available include:

- ◆ **Nifedipine immediate-release only** – 5-10 mg PO or bitten then swallowed. Repeat q 30 minutes until desired outcome achieved. Max dose 30 mg.
- ◆ **Labetalol oral** – 200 mg PO, repeat q 1 hour if needed to achieve goal Max 1200 mg in 24 hours.
- ◆ **Labetalol IV** – 10 mg IV, double dose q 10 minutes to 80 mg until the desired outcome is achieved. Max dose 300 mg, then switch to oral (do not use if congestive heart failure, asthma, or hypovolemic shock).

Referral

- **Organize reliable transportation**
- **Communicate with receiving facility**
 - Explain her diagnosis and condition
 - Describe care provided including medication
 - Estimate her time of arrival
- **Accompany by a provider and companion**
 - Monitor fetus/newborn
 - Maintenance dose
 - IV access maintained



Discuss: What are the steps in executing an effective referral? Guide participants with the following responses:

- ◆ Organize reliable transportation.
- ◆ Communicate with the referral facility:
 - ❖ Explain her diagnosis and condition.
 - ❖ Describe care provided, including medicines.
 - ❖ Estimate her time of arrival.
- ◆ Ensure that the woman is accompanied by a companion.
- ◆ If possible, send a provider to go with her to:
 - ❖ Monitor her and the fetus/newborn.
 - ❖ Provide the first maintenance dose if she does not reach the referral facility within 4 hours.
 - ❖ Keep her IV in place.
 - ❖ Complete the referral record to send with her.
- ◆ Record the referral.
- ◆ Obtain a report from the referral facility and ensure that she receives follow-up care.

Unit 8: Rapid Assessment and Management

Time

45 minutes

Objectives

By the end of this unit, participants will be able to:

- ◆ Quickly identify and treat an obstetric emergency.
- ◆ Treat symptoms of shock.

Materials

- ◆ Presentation
- ◆ Normal vaginal delivery supplies
- ◆ IV fluids
- ◆ Simulated medications

Instructions

This unit will discuss rapid assessment and management of a patient presenting with an unknown obstetric emergency. It is important to perform a rapid assessment of the patient at the first meeting to quickly identify conditions requiring referral. Initial rapid assessment and management when a patient arrives will help you decide whether you are able to treat her at the health center level or if she needs immediate referral to a higher level of care.

Ask the participants: What are some signs of an obstetric emergency? Possible answers may include the below:

- ◆ Unconscious
- ◆ Convulsing
- ◆ Headache
- ◆ Visual disturbances
- ◆ Bleeding
- ◆ Difficulty breathing
- ◆ Fever
- ◆ Severe vomiting
- ◆ Severe abdominal pain

Please note that the above list is not a comprehensive list of obstetric emergencies.

Walk the participants through the identification and treatment of shock using the **slide presentation**. Explain that this type of obstetric emergency must be quickly identified, appropriate treatment rapidly initiated, and referral completed as soon as possible. Discuss the signs of shock including rapid pulse, low blood pressure, pallor, and sweating.

Ongoing practice is key for quickly identifying and treating these emergencies. Patients with shock may not present frequently in a care setting, but when they do, the acuity of their condition is high enough to warrant preparedness.

Presentation



Unit 8: Initial rapid assessment and management

Basic emergency obstetric and newborn care
in humanitarian settings



Rapid assessment and management (RAM)

- Assess the general condition of the woman immediately on arrival
- Quickly identify an emergency
- Prepare to rapidly treat and refer to a higher level of care, as needed

Pregnancy, childbirth, postpartum and newborn care: a guide for essential practice – 3rd ed.



Your ability to provide appropriate care for a woman with an obstetric emergency will depend on the resources available at the facility where you are working. Rapid referral to a higher level of care may be needed. It is important to perform a rapid assessment of the patient at the first meeting to quickly identify conditions requiring referral. An initial rapid assessment when a patient arrives will help you decide whether you are able to treat her at the health center level or if she needs immediate referral to a higher level of care.

Ask: What are some signs of an obstetric emergency? Possible answers are listed below.

- ◆ Unconscious
- ◆ Convulsing
- ◆ Headache
- ◆ Visual disturbances
- ◆ Bleeding
- ◆ Difficulty breathing
- ◆ Fever
- ◆ Severe vomiting
- ◆ Severe abdominal pain

A woman presents to the maternity ward:

Check:

1. Is she in shock?
2. Is there vaginal bleeding?
3. Are there convulsions? Is she unconscious?
4. Is there severe abdominal pain?
5. Is there a dangerous fever?
6. Is she having difficulty breathing?
7. Are there other danger signs present?

Systematically assess a woman of childbearing age for any of the emergencies that we discussed thus far in the training. Read through the questions. **Ask the participants:** What are the symptoms of shock?

Signs of Shock

Early shock	Late shock
Awake, aware, anxious	Confused or unconscious
Fast pulse	Very fast and weak pulse, rate of 110 per minute or more
Slightly fast breathing	Fast and shallow breathing (rate of 30 per minute or more)
Pale	Marked pallor, especially of inner eyelid, palms or around mouth
Sweatiness	Cold, clammy skin
Low blood pressure	Very low blood pressure
Urine output of 30ml per hour or more	Urine output of less than 30ml per hour

Ref Midwifery Education Modules, WHO

Review the signs of shock. Explain that low blood pressure is defined as a systolic blood pressure less than 90 mmHg.

Treat shock

- Put the patient on her left side with her legs elevated
- Insert IV line and start fluids – rapid rate
- Cover with a blanket for warmth
- If no IV access – oral rehydration solution
- Initiate hospital transfer



Shock requires immediate treatment followed by further assessment to identify the rest of the clinical picture of the patient. Call for help immediately.

Insert an IV line and start an IV infusion at a rapid rate. This patient will need to go to the hospital.

Stay or go?

- Speed is crucial when handling emergencies
- Work together to start emergency care and initiate the transfer process
- Know the resources at your facility
- Know the referral process and to where the patient can go for a higher level of care



Unit 9: Newborn Resuscitation (Optional)

This unit was extracted from Unit 4, Practical Session C in the Assisted Vaginal Delivery via Vacuum Extraction clinical refresher training module by the IAWG Training Partnership Initiative.

Time

45 minutes

Objectives

By the end of this unit, participants will be able to:

- ◆ Practice newborn resuscitation using a bag and a mask.

Materials

- ◆ Newborn simulator(s) x 2
- ◆ Neonatal bag and mask
- ◆ Mucus extractor or other type of neonatal suction apparatus
- ◆ Fabric or towel to dry and wrap the baby
- ◆ Clean delivery supplies
- ◆ *Checklist for Newborn Resuscitation* handout, one per participant

Instructions

- ◆ Prepare two identical work stations in one room with the newborn simulators and all necessary equipment as listed above. Place ample space between the two demonstration sites.
- ◆ Each participant should perform the entire process of newborn resuscitation with a baby simulator at least once during the practical session.
- ◆ Participants should take turns using a checklist to check the skills of a peer performing the simulation.
- ◆ The trainer supervises the session and provides feedback to the group.
- ◆ Give a demonstration prior to letting the participants practice.

Presentation



Unit 9: Newborn Resuscitation (Optional)

Basic emergency obstetric and newborn care in humanitarian settings



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISIS

CHECKLIST FOR NEWBORN RESUSCITATION (Many of the following steps/tasks should be performed simultaneously)				
STEP/TASK			REMARKS	
Getting Ready (Prepare for a birth)				
1. Ensure that the area for newborn resuscitation is prepared and that a mucus extractor, self-inflating bag, correct-sized masks for ventilation, and pediatric stethoscope are clean and ready to use for every delivery. Provider should have washed hands and put on sterile gloves.				
2. Tell the woman (and her support person) what is going to be done and encourage them to ask questions.				
3. Provide continual emotional support and reassurance as feasible.				
Skill/Activity Performed Satisfactorily?	Initials		Date	
Immediate Newborn Care				
1. When a baby is born, place immediately on mother's abdomen and dry the baby quickly and thoroughly with a warm, dry cloth.				
2. Assess baby's crying and breath effort during the first 60 seconds after birth. If crying/breathing normally continue with the next step. If not crying or breathing normally by 60 seconds after delivery, go to "Initial Resuscitation Step #1."				
3. Remove wet cloth and place baby skin-to-skin on mother's chest, covering with a warm, dry cloth. Cover head with cap or cloth.				
4. Clamp and cut cord within 2-3 minutes or after pulsations have ceased.				
5. Continue to observe baby's breathing/crying as you proceed with other steps of the delivery.				
Skill/Activity Performed Satisfactorily?	Initials		Date	
Initial Resuscitation Steps (if the baby does not cry or not breathing normally)				
1. Dry the baby quickly and thoroughly. Remove the wet cloth.				
2. Clear the airway; position head and suction mouth and nose only if secretions are seen. Do not suction mouth and nose routinely.				
3. Stimulate breathing by rubbing back 2-3 times.				
4. If the baby cries or breathes normally, place baby skin-to-skin on mother's chest, covering with a warm, dry cloth. Cover head with cap or cloth.				
5. If the baby does not breathe after rubbing the back, clamp and cut the cord, place the baby on a clean, dry surface in the resuscitation area, and cover with a hat and dry cloth, leaving the chest exposed.				
6. Proceed with ventilation using bag and mask within one minute after birth.				
Skill/Activity Performed Satisfactorily?	Initials		Date	

CHECKLIST FOR NEWBORN RESUSCITATION (Many of the following steps/tasks should be performed simultaneously)				
STEP/TASK			REMARKS	
Resuscitation using bag and mask				
1. Position the baby's head in a slightly extended position to open the airway.				
2. Place the mask on the baby's face so that it covers the chin, mouth and nose. Form a seal between mask and face and begin ventilation.				
3. Ensure that the chest is rising with each ventilation. Ventilate at a rate of 40 breaths/minute for 1 minute.				
4. If the baby is still not breathing, call for help and improve ventilation. ▼ Head – reposition, reapply mask ▼ Mouth – clear secretions, open mouth slightly ▼ Bag – squeeze harder and continue ventilation				
5. If not breathing well, palpate umbilical cord or listen to hear rate with stethoscope. ▼ If heart rate is more than 100, continue ventilation. If the baby is breathing spontaneously and there is no in-drawing of the chest and no grunting, put the baby in skin-to-skin contact with the mother. ▼ Monitor with mother.				
6. If breathing is less than 30 per minute, heart rate is less than 100 per minute or severe chest in-drawing is present continue ventilating (with oxygen is available) and arrange for immediate referral for advanced care.				
7. If the baby does not breath spontaneously and has no detectable heart rate after 10 minutes of ventilation, resuscitation should be stopped.				
8. If the baby has a heart rate below 60 beats per minute and no spontaneous breathing after 20 minutes of ventilation, resuscitation should be stopped.				
Skill/Activity Performed Satisfactorily?	Initials		Date	
Post-procedure tasks				
1. Place disposable suction catheters and mucus extractors in a leak-proof container or plastic bag. Place reusable catheters and mucus extractors in 0.5% chlorine solution for decontamination. Then clean and process.				
2. Clean and decontaminate the valve and mask and check for damage.				
3. Wash hands thoroughly.				
4. Ensure that the mother is aware of the outcome of the resuscitation and provide support as necessary.				
5. Record pertinent information on the mother's/newborn's record.				
Skill/Activity Performed Satisfactorily?	Initials		Date	

Prepared by Jhpiego for Ministry of health in the Republic of the Union of Myanmar. (2012).

Improved Midwifery for Maternal, Newborn and Child Health Services: Best Practices in Maternal and Newborn Care Facilitator's Manual. Used with permission.

Unit 10: Partogram (Optional)

This unit was extracted from Unit 4, Practical Session E in the Assisted Vaginal Delivery via Vacuum Extraction clinical refresher training module by the IAWG Training Partnership Initiative.

Time

45 minutes

Objectives

By the end of this unit, participants will be able to:

- ◆ Use the partogram to chart labor progress.

Materials

- ◆ *Partogram*, three copies of a blank handout per participant
- ◆ *Partogram - answers*, one copy of handout per participant
- ◆ Video (on the USB key and IAWG website)

Instructions


This session is optional. It is recommended that the cases are given to the participants during Unit 9 (also optional) so they can work on them when they are not participating in the newborn resuscitation activity. Make time to debrief on this activity, if it is included. Instruct the participants to complete the partogram in groups of 2 or 3. Each case study provides instructions for charting information on the partogram and making clinical decisions based on the information provided. The answer key contains pictures of completed partograms and corresponding case study notes for discussion. Participants should receive the answer key after the activity is complete.

Note to Trainers: If you opt to do this session, review the partogram form with the group before beginning the exercise. You can do so using the slides in this unit and the 16-minute partogram video, based on participants' familiarity with the partogram.

Presentation



Partogram




IAWG

INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISES

Give a brief overview of the partogram using this and the following slide.

Note: There is a link to a video about the partogram (16 minutes) included in the training package. Watching the video and doing the associated exercises (see the IAWG website or USB key) is at the discretion of the trainer, and based on the needs of the participants and time.

Partogram



IAWG

INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISES

The partogram (or partograph) is a structured, graphical representation of the progress of labor. It shows cervical dilatation in relation to time as well as the descent of the fetal head. Space can be provided to write notes on the partogram including: the frequency of contractions, fetal heart rate, maternal observations, and medication administration. By using it, the progress of labor can be seen at a glance on one sheet of paper. The partogram is a tool to identify when intervention is necessary, as failure to progress can be easily recognized.

Unsatisfactory progress in labor and the possibility of intervention is assessed when cervical dilatation is too far to the right of the alert line on the partogram.

Practical Session Answer Key

Adapted from Jhpiego. *Managing Complications in Pregnancy and Childbirth Learning Resource Package*.

INSTRUCTIONS: There are three case studies about labor management. Follow each step of the assigned case and fill out the partogram as you go along, plotting the data as it is provided. Answer the associated questions. Discuss the questions with your team as you go along. Review the answer key for any discrepancies with your exercise. These will be discussed as a group at the end of the session.

CASE 1

INSTRUCTIONS: The group members should systematically go through each step below and plot the information on their individual partograms. Answer the questions associated with each step for Mrs. A.

STEP 1: Mrs. A was admitted at 09.00 on 19 September 2015. Membranes ruptured at 08.00. She is a gravida 3, para 2+0. The hospital number is 7886. On admission, the fetal head was 4/5 palpable above the symphysis pubis and the cervix was 2 cm dilated. Contractions are 2 in 10, lasting 25 seconds.

Q: What should be recorded on the partogram?

Note: Mrs. A is not in the active phase of labor. Record the details of her history, contractions, and your examination findings. If you are using a partogram that allows for observations of the latent phase of labor, record your findings on the partogram. If not, record your findings in the mother's case notes. She is due for another vaginal examination in 4 hours.

SEE PARTOGRAM

STEP 2: It is 13.00 (4 hours later). The fetal head is 2/5 palpable above the symphysis pubis. The cervix is 5 cm dilated.

Q: What should you now record on the partogram?

SEE PARTOGRAM

STEP 3: Labor is progressing. Mrs. A is now in the active phase of labor. Note the phase of labor and plot the following information on the partogram:

- The fetal head is 2/5 palpable above the symphysis pubis
- The cervix is 5cm dilated
- 3 contractions in 10 minutes, each lasting 40 seconds
- Fetal heart rate (FHR) 120
- Membranes ruptured, amniotic fluid clear on pad check
- Sutures of the skull bones are apposed (molding +), no caput
- Blood pressure 120/70 mmHg

- Temperature 36.8°C
- Pulse 80/minute
- Urine output 150 mL; negative protein and acetone

Answer the following questions:

Q: What steps should be taken?

- Inform Mrs. A and her family of the findings and what to expect.
- Encourage Mrs. A to ask questions.
- Provide comfort measures and hydration.

Q: What advice should be given?

- Assume the position of her choice.
- Drink plenty of fluids.
- Eat as desired.

Q: What do you expect to find at 17:00?

- Progress to at least 9 cm dilatation.

STEP 4: Labor monitoring is ongoing over the next four hours. Plot the following information on the partogram:

13.30	FHR 120, Contractions 2/10 each 30 seconds, Pulse 80/minute
14.00	FHR 136, Contractions 2/10 each 30 seconds, Pulse 80/minute
14.30	FHR 140, Contractions 2/10 each 35 seconds, Pulse 88/minute
15.00	FHR 130, Contractions 2/10 each 40 seconds, Pulse 88/minute
15.30	FHR 136, Contractions 3/10 each 40 seconds, Pulse 84/minute
16.00	FHR 140, Contractions 3/10 each 40 seconds, Pulse 88/minute
16.30	FHR 130, Contractions 3/10 each 45 seconds, Pulse 88/minute
17.00	FHR 140, Contractions 3/10 each 45 seconds, Pulse 90/minute

Answer the following questions:

Q: What steps should be taken?

- Assume the position of her choice.
- Frequent sips of fluids.

Vaginal assessment at 17.00 shows that the cervix is now fully dilated and the head has descended to 0/5; Mrs. A now feels expulsive. Record the following information on the partogram.

- FHR 140, contractions 4/10 each lasting 50 seconds, pulse 80/minute, temperature 38 C

Q: What advice should be given?

- Push only with the urge to push.

Q: What do you expect to happen next?

- Spontaneous vaginal birth.

STEP 5: Record the following information on the partogram:

17.20 Spontaneous birth of a live female infant weighing 2,850 g

Answer the following questions:

Q: How long was the active phase of the first stage of labor?


- 5 hours (4 hours from 13.00 - 17.00).

Q: How long was the second stage of labor?

- 20 minutes.

CASE 1

Name Mrs. A Gravida 3 Para 2 +0 Hospital number 7886 (464521)
 Date of admission 19.9.15 Time of admission 9 AM Ruptured membranes 1 hours



CASE 2

INSTRUCTIONS: The group should systematically go through each step below and plot the information on each participant's individual partogram. Answer the questions associated with each step for Mrs. B.

STEP 1: Mrs. B was admitted at 10.00 on 19 September 2013. She is gravida 1, para 0+0. Her membranes are intact. The hospital number is 1443.

Record the personal history details above on the partogram, together with the following details:

- The fetal head is 3/5 palpable above the symphysis pubis
- The cervix is 4 cm dilated
- 2 contractions in 10 minutes, each lasting for 30 seconds
- FHR 140
- Membranes intact
- Blood pressure 100/70 mmHg
- Temperature 36.2°C
- Pulse 80/minute

Answer the following questions:

Q: What is your diagnosis?

- Active labor

Q: What action will you take?

- Inform Mrs. B and her family of the findings and what to expect.
- Always provide the opportunity for Mrs. B to ask questions.
- Encourage Mrs. B to walk around, eat, and drink as desired.

STEP 2: Plot the following information on the partogram for Mrs. B:

10.30	FHR 140, Contractions 2/10 each 25 seconds, Pulse 90/minute
11.00	FHR 136, Contractions 2/10 each 15 seconds, Pulse 88/minute
11.30	FHR 140, Contractions 2/10 each 30 seconds, Pulse 84/minute
12.00	FHR 136, Contractions 2/10 each 30 seconds, Pulse 88/minute
12.30	FRH 136, Contractions 2/10 each 30 seconds, Pulse 90/minute
13.00	FRH 140, Contractions 2/10 each 30 seconds, Pulse 88/minute
13.30	FRH 130, Contractions 2/10 each 30 seconds, Pulse 88/minute
14.00	FRH 140, Contractions 2/10 each 30 seconds, Pulse 90/minute

- Temperature 36.2 C
- Blood Pressure 100/70 mmHg
- Urine output 300 mL: negative protein and acetone
- The cervix is 4 cm dilated
- The fetal head is 3/5 palpable above the symphysis pubis
- Membranes intact

Answer the following questions:

Q: What is your diagnosis?

- Prolonged active phase of labor.
- Less than 3 contractions in 10 minutes, each lasting less than 40 seconds.
- Good fetal and maternal condition.

Q: What action will you take?

NOTE: Plan to facilitate a discussion about using oxytocin to augment labor based on the clinical setting. For instance, is the woman being cared for at a health post that is 4 hours away from a district hospital where an oxytocin drip can be started? Or if she is in a district hospital, can other measures be used before oxytocin is started, such as hydration and ambulation?

Q: What is your diagnosis?

- Prolonged active phase.
- Less than 3 contractions per 10 minutes, each lasting less than 30 seconds.
- Good fetal and maternal condition.

Q: What will you do?

- Augment labor with oxytocin and artificial rupture of membranes.
- Inform Mrs. B and her family of the findings and what to expect.
- Reassure.
- Answer questions.
- Encourage drinks.
- Encourage Mrs. B to assume the position of her choice.

STEP 3: Plot the following information on the partogram:

- | | |
|-------|--|
| 14:00 | <ul style="list-style-type: none">– The cervix is 4 cm dilated, sutures apposed– Membranes artificially ruptured, clear fluid– Labor augmented with oxytocin 2.5 units in 500 mL IV fluid at 10 drops per minute (dpm) |
| 14.30 | <ul style="list-style-type: none">– 2 contractions in 10 minutes, each lasting 35 seconds– FHR 140, Pulse 90/minute– Infusion rate increased to 20 dpm |
| 15.00 | <ul style="list-style-type: none">– 3 contractions in 10 minutes, each lasting 40 seconds– FHR 140, Pulse 90/minute– Infusion rate increased to 30 dpm |
| 15:30 | <ul style="list-style-type: none">– 3 contractions in 10 minutes, each lasting 45 seconds– FHR 140, Pulse 88/minute– Infusion rate increased to 40 dpm |
| 16.00 | <ul style="list-style-type: none">– Fetal head now 2/5 palpable above the symphysis pubis– Cervix 6 cm dilated; sutures apposed– 3 contractions in 10 minutes, each lasting 45 seconds– FHR 144, Pulse 92/minute– Amniotic fluid clear– Infusion rate maintained at 40 dpm |

- 16.30 – 3 contractions in 10 minutes, each lasting 45 seconds
 – FHR 140, Pulse 90/minute
 – **Infusion increase to 50 dpm**

Q: What steps would you take?

- Continue to augment labor (maintain oxytocin infusion rate at 50 dpm).
- Provide comfort (psychological and physical).
- Encourage drinks and nutrition.

STEP 4: Plot the following information on the partogram:

- 17.00 FHR 138, Pulse 92/minute, Contractions 3/10 each 45 seconds, Maintain at 50 dpm
17.30 FHR 140, Pulse 94/minute, Contractions 3/10 each 45 seconds, Maintain at 50 dpm
18.00 FHR 140, Pulse 96/minute, Contractions 4/10 each 50 seconds, Maintain at 50 dpm
18.30 FHR 144, Pulse 94/minute, Contractions 4/10 each 50 seconds, Maintain at 50 dpm

STEP 5: Plot the following information on the partogram:

- 19.00 – Fetal head 0/5 palpable above the symphysis pubis
 – 4 contractions in 10 minutes, each lasting 50 seconds
 – FHR 144, Pulse 90/minute
 – Cervix fully dilated. Mother feels expulsive.

STEP 6 : Record the following information on the partogram:

- 19.30 – 4 contractions in 10 minutes, each lasting 50 seconds
 – FHR 142, Pulse 100/minute
- 20.00 – 4 contractions in 10 minutes, each lasting 50 seconds
 – FHR 146, Pulse 110/minute
 – Spontaneous birth of a live male infant weighing 2,654 g

Answer the following questions:

Q: How long was the active phase of the first stage of labor?

Nine hours.

Q: How long was the second stage of labor?


One hour (60 minutes).

Q: Why was labor augmented?

Less than 3 contractions in 10 minutes, each lasting less than 40 seconds (lack of progress).

CASE 2

Name Mrs. B Gravida 1 Para 0⁺⁰ Hospital number 1443
 Date of admission 19.9.13 Time of admission 10 AM Ruptured membranes 1400 hours



CASE 3

STEP 1: Mrs. C was admitted at 10.00 on 19 September 2013. Membranes ruptured spontaneously at 04.00. She is a gravida 4, para 3+0. Her hospital number is 6639.

Record the information above on the partogram, together with the following details:

- Fetal head 3/5 palpable above the symphysis pubis Cervix 4 cm dilated
- 3 contractions in 10 minutes, each lasting 30 seconds FHR 140
- Amniotic fluid clear
- Sutures apposed (Molding +)
- Blood pressure 120/70 mmHg
- Temperature 36.8°C
- Pulse 80/minute

STEP 2: Plot the following information in the partogram:

10.30 FHR 130, Contractions 3/10 each 45 seconds, Pulse 80/minute
11.00 FHR 136, Contractions 3/10 each 45 seconds, Pulse 90/minute
11.30 FHR 140, Contractions 3/10 each 45 seconds, Pulse 88/minute
12.00 FHR 140, Contractions 3/10 each 45 seconds, Pulse 90/minute, Temperature 37°C, Head 3/5 palpable
12.30 FHR 130, Contractions 3/10 each 45 seconds, Pulse 90/minute
13.00 FHR 130, Contractions 3/10 each 50 seconds, Pulse 88/minute
13.30 FHR 120, Contractions 3/10 each 50 seconds, Pulse 88/minute
14.00 FHR 130, Contractions 3/10 each 50, Pulse 90/minute, Temperature 37°C, Blood Pressure 100/70 mmHg. Fetal head 3/5 palpable above the symphysis pubis. Cervix 6 cm dilated, amniotic fluid clear. Sutures overlapped but reducible (Molding ++).

STEP 3: Plot the following information in the partogram:

14.30 FHR 120, Contractions 3/10 each 45 seconds, Pulse 90/minute, Clear fluid
15.00 FHR 120, Contractions 3/10 each 45 seconds, Pulse 88/minute, Blood-stained fluid
15.30 FHR 100, Contractions 3/10 each 45 seconds, Pulse 100/minute
16.00 FHR 90, Contractions 3/10 each 50 seconds, Pulse 100/minute, Temperature 37°C
16.30 FHR 96, Contractions 4/10 each 50 seconds, Pulse 110/minute. Fetal head 3/5 palpable above the symphysis pubis. Cervix 6 cm dilated. Amniotic fluid meconium stained. Sutures overlapped and not reducible. Urine output 100 mL; protein negative, acetone 1+.

STEP 4: Record the following information on the partogram:

Cesarean section at 17.30, live female infant with poor respiratory effort and weighing 4,850 g.

Answer the following questions:

Q: What is the final diagnosis?

Obstructed labor with fetal head 3/5 palpable above the symphysis pubis.

Q: What action was indicated at 14.00, and why?

Cesarean section (CS) because Mrs. C was already in secondary arrest of dilatation and descent despite at least 3 contractions in 10 minutes, each lasting more than 40 seconds.

Q: What action was indicated at 15.00, and why?

- Continue emotional and physical support, including hydration.
- Continue attentive monitoring of maternal and fetal condition.
- Have crossed the alert line.
- Blood-stained amniotic fluid.

Q: At 17.00, a decision was taken to do a cesarean section, and this was rapidly done. Was this a correct action?


Yes, it was correct because of the following: the fetal condition was deteriorating, failure to progress, and a rising maternal pulse.

Q: What problems may be expected in the newborn?

Asphyxia, meconium aspiration.

CASE 3

Name Mrs. C Gravida 4 Para 3⁺⁰ Hospital number 6639
 Date of admission 19.9.13 Time of admission 10 AM Ruptured membranes 4 AM hours



Unit 11: Closing

Time

30 minutes

Objectives

By the end of this unit, participants should be able to:

- ◆ Explain how the training met their expectations and course objectives.


Materials

- ◆ Flip chart paper
- ◆ *Course Evaluation*, one per participant
- ◆ *Post-test*, one per participant
- ◆ *Certificate of Completion*, one per participant

Instructions

End the training by first reviewing the course objectives and revisiting the participants' expectations. Ensure that all expectations have been addressed. Hand out the *Post-test* and ask the participants to take 15 minutes to complete it. After finishing the test, participants can take 10 minutes to fill out the *Course Evaluation*. Thank the participants for their attention and participation in the short refresher training course. Present each participant with a prepared *Certificate of Completion* to wrap up the training.

Presentation



Sample Certificate of Attendance

Certificate of Completion

[Recipient Name]

is hereby recognized as completing a two-day clinical refresher course on

**Basic Emergency Obstetric and
Newborn Care in Humanitarian Settings,
Select Signal Functions**



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISES

PRESENTED BY:

ON THIS DAY:

References

- Jhpiego. *Helping Mothers Survive – Bleeding after birth*. 2015.
- Jhpiego. *Helping Mothers Survive – Pre-eclampsia/eclampsia*. 2015.
- Médecins Sans Frontières. *Essential Obstetric and Newborn Care*. 2015 edition.
- Medical Aid Films. *How to manage bleeding after birth*.
- Medical Aid Films. *How to perform manual removal of the placenta*.
- White Ribbon Alliance. *Respectful maternity care: The universal rights of childbearing women*. 2011.
- World Health Organization. *Recommendations for Prevention and Treatment of Maternal Peripartum Infections*. 2015.
- World Health Organization. *Pregnancy, childbirth, postpartum and newborn care – An essential guide*. 3rd Ed. 2015.
- World Health Organization. *Education material for teachers of midwifery: midwifery education modules*. 2nd ed. *Managing postpartum hemorrhage*. 2008.
- World Health Organization. *Education material for teachers of midwifery: midwifery education modules*. 2nd ed. *Managing eclampsia*. 2008.
- World Health Organization. *Education material for teachers of midwifery: midwifery education modules*. 2nd ed. *Managing puerperal sepsis*. 2008.
- United Nations Population Bureau. *Inter-agency Reproductive Health Kits for Crisis Situations*. 5th ed. 2011.



INTER-AGENCY WORKING GROUP
ON REPRODUCTIVE HEALTH IN CRISES



USAID
FROM THE AMERICAN PEOPLE